

Evaluation and Correlation of Heparin-25 Hormone, Hemoglobin and Packed Cell Volume of Patients with β -Thalassaemia Intermedia Before and After Blood Transfusion for Iraqi Children Patients

Mohammed Q. saadon

Alasheqi@yahoo.com

Bushra H. Ali

Drbushra750@yahoo.com

Department of Chemistry, College of Education for Pure Science Ibn Al-Haitham, University of Baghdad, Baghdad, Iraq.

Article history: Received 26 June 2018, Accepted 10 July 2018, Published December 2018.

Abstract

β -thalassaemia consists of 3 chief forms: thalassaemia primary (other called "cooley's Anaemia" or "Mediterranean durability Anaemia"), thalassaemia intermedia or thalassaemia minor also frequent termed " β -thalassaemia carrier", " β -thalassaemia trait" however, "heterozygous β -thalassaemia". separately from the rare magisterial forms, problem along β -thalassaemia essential are homozygotes or made heterozygotes because B^0 and B^+ genes, problem including thalassaemia intermedia are typically homozygotes and compound heterozygotes then subjected with thalassaemia minor are broadly heterozygotes.

In this study, we take eighty humans the age about them had been mean \pm SD (9.68 \pm 2.08), 40 patients along β -thalassaemia intermedia, then 40 healthy people as much control. extraction gore 5ml from thalassaemia's patients before gore transfusion then afterwards take the blood by way of three days. The samples we instituted are 120, forty samples before blood transfusion, 40 samples after gore transfusion then 40 control samples. Then we separated every pattern of twin's tubes.

Amplify blood attention about β -thalassaemia intermedia patient since blood advancement due according to minimize degree regarding hepcidin-25

further than its instruction was once shown up iron focus meanly affect through hepcidin-25 level, therefore so hepcidin-25 used to be surprisingly significant exchange in β -thalassaemia intermedia affected person afterwards blood advancement yet so much conduct to iron overload. So we conclude low level of Hemoglobin before and after blood transfusion due to decrease or absent formation of β -globin that because gene disorder, and conclude correlation between hepcidin hormone with Hb and PCV.

The aim of this study is to evaluate hepcidin-25 levels, hemoglobin [Hb] and packed cell volume [PCV] before and after blood transfusion, then correlation of hepcidin-25 hormone with [Hb] and [PCV] of children patients with β -thalassaemia intermedia.

Keywords: β -thalassaemia intermedia, Hemoglobin and hepcidin-25.

1. Introduction

Thalassemia refers to conformity with a cluster regarding gore ailments stability characterized via mean or poor introduction concerning normal globin manacles [1]. According to the series whose creation is harmed durability such namely α -, β -, γ -, σ - and

$\epsilon\gamma\beta$. Mainly severe thalassemia is followed at the equal period so recessive toughness behavior. These imperative toughness quantitative defects are no longer hermetically separated of the structural alternates built through the facet over condensed durability dimension [such as much HbE durability toughness than Hb Lepore]. Starting toughness permanency, a scientific toughness factor regarding observation, the most types are α - and β -thalassemia, reduce certain over the pair kinds about polypeptide manacles [α then β] as form everyday hemoglobin particle [HbA, $\alpha_2\beta_2$] [2].

Therefore, Patients with β -thalassemia intermedia, important modifier of phenotype is the wide change of mutations that have an effect on the β -globin gene among the homozygous then made-up heterozygous state [3]. longevity permanency The length beside average farmer mutations fabulous between consequence on slight limit on β -globin puttie production in accordance after the deep different mutations as stop result among the β_0 -thalassaemias; upon in accordance with expectation is, a completed non-appearance concerning β -globin bandage synthesis. Deletions over the β -globin gene are rare [4]. The measure regarding mutations but consonant changeable dimension over α/β -globin strip imbalance but insensitive erthropoiesis are the essential decided due to the fact milder anaemia then phenotype between β -thalassemia intermedia than β -thalassemia fundamental [5]. Secondary modifiers are involved at once of modifying the content above α/β -globin tuck imbalance encyclopaedic coinheritance over brilliant molecular forms concerning α -thalassemia, prolonged problem concerning α -hemoglobin stabilizing proton, yet nice concord on γ -chains among adult lifestyles [6]. Several genes undergo commence as like function according to modify γ -chain technical yet honor phenotype, some touching in imitation of them encoded among the β -globin gene cluster, others are amongst special chromosomes [7]. While tertiary modifiers incorporate polymorphisms on in accordance with hope are no longer related in accordance according to globin puttie technical however may additionally moreover keep an enhance have an effect on related to unique issues over the sickness so (iron absorption, bilirubin metabolism, skeleton metabolism, cardiovascular disease, and susceptibility within imitation along infection)[8] Therefore, β -thalassemia intermedia may additionally moreover additionally end result past outcome industrial about α -globin chains with the aid of triplicated and quadruplicated α -genotype associated along with β -heterozygosity [9]. lifetime Blood Less commonly, a alone β -globin place affect, the vile lifestyles totally normal, subsequently β -thalassemia intermedia is dominantly traditional [10].

Therefore Oxygen was once received in imitation of the tissues from the courage by using the enormously specialised protein particle, hemoglobin intimate the purple compartments concerning the blood, so was dwell placed all pink gore compartment holds about 300 bags of molecules of that protein, totally a propos 30 picograms of matter because of each mobilephone [11]. Everyone bit of hemoglobin used to be built via two pairs concerning matching sub-unit; globin manacles are named along the slip concerning the Greek alphabet and match of pair groups: the α -globin paint brush consists the ζ - afterward α -globin chains, but the β -globin cluster, component the globin chains ϵ , γ , β yet δ toughness [12].

In fact, the globin chains show off sequentially at partial board in ontogeny and, beneath coupling, the consonant 4 majors' kinds above hemoglobin:

1. "embryonic" hemoglobin's, used to be once best detectable beyond the third since the tenth sennight on gestation and show on $\zeta_2\varepsilon_2$ [Hb Gower 1], $\alpha_2\varepsilon_2$ [Hb Gower 2], $\zeta_2\gamma_2$ [Hb Portland 1]; or $\zeta_2\beta_2$ tetramers [Hb Portland 2];
2. "foetal" hemoglobin [Hb F], constitutes the lordly oxygen company at some flooring within pregnancy or is $\alpha_2\gamma_2$ molecule;
3. "adult" hemoglobin [Hb A $\alpha_2\beta_2$], choice Hb F swiftly under birth;
4. A baby adult component, Hb A2 [$\alpha_2\delta_2$].

The specific hemoglobin kind were constructed but commend above at assured period regarding racial improvement is referred to specifically "hemoglobin switching"[13]. The red gore cells concerning the grown-up endemic correspond about 97-98% regarding Hb A, 2-3% respecting Hb A2 then traces over Hb F [14].

In adult life, the globin adjustment take place between the erythroblast into the bone marrow [15]. Globin chains have the restore shape yet keep association in such an access up to expectation the wide variety about α -chains should exactly healthy to that amount on the β -chains [16].

In summation Hepcidin-25 is a protein among people is encode including the resource about the hepcidin-25 antimicrobial peptide [HAMP] gene, is a resolution director on the entry on blood within the foreign money about mammals. When hepcidin-25 dosage is abnormally extending inflammation, serum metallic shower due according in accordance with iron trapping inside macrophages and mettle cells afterward vile intestine blood absorption, that government within imitation along anemia fit to no longer enough aggregate concerning serum metal lifestyles reachable due to the fact growing crimson cells [17]. However, so the hepcidin-25 stage is abnormally vile as like on hemochromatosis, gore overload shows over suit between imitation along accelerated ferroportin mediated blood efflux from storage but extended intestine metal absorption [18].

So hepcidin-25, a peptide hormone is typically coordination inside the liver, was once located among 2000. It remains capable remain reduces extracellular metal within the body via endless mechanisms:

- 1) Hepcidin-25 decrease dietary metal attention by way of decreasing metallic put all through gut mucosal cells (enterocytes); It reduces metal emerged beyond macrophages, the imperative web page in regard to iron storage;
- 2) It reduces metal backyard in company of the liver. Therefore, into all 3 instances as is accomplice via capacity about reducing the transmembrane iron transporter ferroportin.

Durability longevity toughness permanency longevity durability permanency Hepcidin-25 determined namely a preprohormone (84 amino acids), prohormone (60 amino acids), after hormone (25 amino acids). Twenty- afterwards 22-amino lotos reckless metabolites of hepcidin-25 also located amongst the urine. Removal respecting 5 N-terminal amino acids consequences deactivation regarding feature [19]. The changing regarding prohepcidin according according to hepcidin-25 is intercede through the prohormone convertase furin. This variation can also moreover continue to be alter via alpha-1 antitrypsin [20].

Hepcidin-25 does stand hermetically folded polypeptide along 32% beta blade persona below a hairpin structure stabilized via IV disulfide bonds [21]. In addition, conductor regarding steel metabolism. So hepcidin-25 inhibits gore gait through seize into accordance along the steel export people ferroportin, as like is located regarding the basolateral floor upon gut

enterocytes then the plasma membrane concerning reticuloendothelial cells (macrophages) [22].

2. Material and Method

In this study, we take eighty humans the age about them had been mean \pm SD (9.68 \pm 2.08), 40 patients among them were given β -thalassemia intermedia, while 40 healthful people were used as control sample. We extract gore 5ml from thalassemia's patients before gore transfusion. Afterwards range regarding blood via period 3 days. The volume quantity concerning the samples we done are 120, forty samples before gore transfusion, 40 samples beneath stick length since forty control samples. Then we separated every pattern about pair tubes.

The 1st pipe because of determine Hepsidin-25 and the 2nd longevity barrel according to decide Hemoglobin or PCV. Serum of the samples was separated using centrifuge and the assay by using enzyme-linked immune sorbent assay (ELISA) way to determine Hepsidin-25. The hematological marker like hemoglobin (Hb) and packed cell volume (PCV) was calculated automatically by the hematological tool hemoglobin meter [23].

2.1. Statistical Analysis

Results of this study were expressed as [mean \pm SD], also T.test (spss) was utilized to compare between three studied groups, so T.test less than or equal of 0.05 was considered significant and highly significant, respectively. Also this study expressed as correlation of hepsidin-25 with all parameter, so correlation value between range (\pm 0.5) was considered significant.

3. Results

The patients were into three groups, the first group [C1] is before blood transfusion, the second group [C2] is after blood transfusion, and the third group [C3] is the control. **Table 1.** showed parameters results of 20th female patients with β -thalassemia intermedia.

Table 1. Results of female patient.

Biometers	[C1] Mean \pm SD	[C2] Mean \pm SD	[C3] Mean \pm SD	Ttest C1VsC2	Ttest C1VsC3	Ttest C2VsC3
Hb g/dL	8.09 \pm 1.57	10.37 \pm 1.34	12.2 \pm 0.41	HS	HS	HS
PCV %	25.02 \pm 4.53	32.1 \pm 4.01	37.62 \pm 1.19	HS	HS	HS
S. Hep pg/mL	13.6 \pm 3.09	11.49 \pm 1.45	25.26 \pm 3.56	NS	HS	HS

4. Discussion

The mean \pm SD of hemoglobin is (BT= 8.09 \pm 1.57) g/dL before blood transfusion which is less than patients after blood transfusion (10.37 \pm 1.34) g/dL; while control patients (12.2 \pm 0.41) g/dL showed higher values after and before blood transfusion as illustrated in **Table 1.**

The results show a significant increase with control ($p \leq 0.01$), which agree with Girard JM study, also Girard JM suggested that red blood cells of thalassemia patients can destroy

peripheral arteries due to gene disorder of globin synthesis that leads to frequent hemoglobin deficiency compared to healthy body [23], which results in a significant increase group 2 (C2) ($p < 0.01$) that caused by blood transfusion of group 2 (C2) patients all that is shown in **Figure 1**. [24].




Figure 1. Hb levels in male, female and control.

In addition, the results show that mean \pm SD of packed cell volume (25.02 ± 4.53) % before blood transfusion gives lower value than same patients after blood transfusion (32.1 ± 4.01) %, whereas the PCV in patients with thalassemia was lower than control (37.62 ± 1.19) % as appeared in **Table 1**.

The results were significantly increased compared with C3 ($p \leq 0.01$), these results agree with Weather all D study who showed the deficiency or absent β -globin is accelerated by sedimentation of erythrocyte [25], gave mean \pm SD of PCV for thalassemia patients after blood transfusion more than C1.

The results of C1 was higher ($p \leq 0.01$) compared to C2, these results agree with Al-Ali HK study, who suggested transfusion of blood increase the sedimentation of red blood cell [26], as shown in **Figure 2**.




Figure 2. PCV levels in male patient, and female patient and control

As well as, the mean \pm SD of Hepcidin-25 (13.6 ± 3.09) pg/mL before and after blood which transfusion (11.49 ± 1.45) pg/mL was less than control (25.26 ± 3.56) pg/mL as appears in **Table 1**.

The results of female patients were insignificant ($p \geq 0.05$) while a significant decrease ($p \leq 0.05$) of T.test between C1 and C2 that approve of Hepcidin-25 secretion affected by blood transfusion as shown in **Figure 3**. and **Table 1**. Hepcidin-25 level decreases after blood transfusion so that increase absorption of iron leading to iron overload, and that approved by Zhao N. study when he was studying matriptase-2. He also noticed an increase in iron absorption with decreasing Hepcidin-25 secretion [27], also Wang Y. study was in agreement with these results when he investigated human hemochromatosis protein mutation and effect of bone morphogentic protein in Hepcidin-25 secretion [28], in addition Rishi G. study was in agreement with these results and displayed effect of transferrin receptor-2 in Hepcidin-25 regulate [29].




Figure 3. Serum Hepcidin-25 levels in male patient, and female patient and control.

Table 2. The correlation of Hepcidin-25 with Hemoglobin and Packed cell volume.

Biometer	Before blood transfusion		After blood transfusion	
	r value	T.test	r value	T.test
Hb	0.009	HS	-0.415	HS
PCV	-0.015	HS	-0.415	HS

Hemoglobin correlation with Hepcidin-25 before blood transfusion product showed non-significant as illustrated in **Figure 4**., while after blood transfusion product revealed highly significant negative correlation.




Figure 4. Correlation between Hep with Hb in female patient.

Therefore, Packed cell volume correlation with Hepcidin-25 before blood transfusion demonstrated non-significant positive correlation, while significant negative correlation observed after blood transfusion as appeared in **Figure 5**.




Figure 5. Correlation between Hep with PCV in male patient.

References

1. Unissa. R.; Monica. B.; Konakanchi. S.; Darak. R.; Keerthana. S. L.; Kumar, S. A. Thalassemia: A Review. *AJPR*. **2018**, *8*, 3, 195-202.
2. Rund. D. Thalassemia: modern medicine battles an ancient disease. *Am J Hematol*. **2016**, *91*, 1, 15-21.
3. Mohanty. D.; Colah. R.B.; Gorakshakar. A.C.; Patel. R.Z.; Master. D.C.; Mahanta. J.; Sharma, S.K. Prevalence of β -thalassemia and other haemoglobinopathies in six cities in India: a multicentre study. *J Community Genet*. **2013**, *4*, 1, 33-42.
4. Cao. A.; Kan. Y.W. The prevention of thalassemia. Cold Spring Harbor perspectives in medicine. *Cold Spring Harb Perspect Med*. **2013**, *3*, 2, 1-15.
5. Camaschella.C.; Nai. A. Ineffective erythropoiesis and regulation of iron status in iron loading anaemias. *Br. J. Haematol*. **2016**, *172*, 4, 512-523.
6. Amoozgar. H.; Zeighami. S.; Haghpanah. S.; Karimi. M. A comparison of heart function and arrhythmia in clinically asymptomatic patients with β thalassemia intermedia and β thalassemia major. *Hematology*. **2017**, *22*, 1, 25-29.

7. Danjou. F.; Francavilla. M.; Anni. F. A genetic score for the prediction of beta-thalassemia severity. *Haematologica*. **2015**, *100*, 4, 452–7.
8. Baldini, M.; Marcon, A.; Cassin, R.; Ulivieri, F.M.; Spinelli, D.; Cappellini, M.D. and Graziadei, G. Beta-thalassaemia intermedia: evaluation of endocrine and bone complications. *Biomed Res Int*. **2014**, *2014*, 1-5, <http://dx.doi.org/10.1155/2014/174581>.
9. Arlet.J.B.; Ribeil. J.A.; Guillem. F.; Negre. O.; Hazoume. A.; Marcion. G. HSP70 sequestration by free α -globin promotes ineffective erythropoiesis in β -thalassaemia. *Nature*. **2014**, *514*, 7521, 242-6.
10. Baronciani, D.; Angelucci, E. and Potschger, U. Hemopoietic stem cell transplantation in thalassemia: A report from the european society for blood and bone marrow transplantation hemoglobinopathy registry,2000-2010. *Biol Blood Marrow Transplant*. **2010**,*51*, 4, 536.
11. Camaschella. C.; Nai. A. Ineffective erythropoiesis and regulation of iron status in iron loading anaemias. *Br. J. Haematol*. **2016**, *172*, 4, 512–523.
12. Thein. S.L. The molecular basis of β -thalassemia. *Cold Spring Harb Perspect Med*.**2013**, *3*, 5,1-24.
13. Taher. A.T.; Musallam. K.M.; Saliba. A.N.; Graziadei. G.; Cappellini. M.D. Hemoglobin level and morbidity in non-transfusion-dependent thalassemia. *Blood Cells Mol. Dis*. **2015**, *55*(2):108–109.
14. Paglietti. M.E.; Satta. S.; Sollaino. M.C.; Barella. S.; Ventrella. A.; Desogus. M.F.; Demartis. F.R.; Manunza. L.; Origa. R. The problem of borderline Hemoglobin A2 levels in the screening for b- thalassemia carriers in Sardinia. *Acta Haematol*. **2016**, *135*(4):193–199.
15. Haase, V.H. Regulation of erythropoiesis by hypoxia-inducible factors. *Blood Rev*. **2013**, *27*, 1, 41-53.
16. Hardison. R.C. Evolution of hemoglobin and its genes. *Cold Spring Harb Perspect Med*. **2012**, *2*, 12, 1-18.
17. Jones.E.; Pasricha. S.R.; Allen. A.; Evans. P.; Fisher. C.A.; Wray. K.; Premawardhena. A.; Bandara. D.; Perera. A.; Webster. C.; Sturges. P. Heparin is suppressed by erythropoiesis in hemoglobin E β -thalassemia and β -thalassemia trait. *Blood*. **2015**, *125*, 5, 873-880.
18. Baronciani. D.; Angelucci. E.; Potschger. U. Hemopoietic stem cell transplantation in thalassemia: A report from the european society for blood and bone marrow transplantation hemoglobinopathy registry, 2000–2010. *Bone Marrow Transpl*. **2016**, *51*, 4, 536–541.
19. Casu. C.; Oikonomidou. P.R.; Chen. H. Minihepcidin peptides as disease modifiers in mice affected by β -thalassemia and polycythemia vera. *Blood*, **2016**, *128*, 2, 265–276.
20. Tarkun. P.; Mehtap. O.; Atesoğlu. E.B.; Geduk. A.; Musul. M.M.; Hacıhanefioglu. A. Serum hepcidin and growth differentiation factor-15 (GDF-15) levels in polycythemia vera and essential thrombocythemia. *Eur J Haematol*. **2013**, *91*, 3, 228-235.
21. Latour. C. Testosterone perturbs systemic iron balance through activation of epidermal growth factor receptor signaling in the liver and repression of hepcidin. *Hepatology*. **2014**, *59*, 2, 683–694.

22. Pasricha. S.R.; Frazer. D.M.; Bowden. D.K.; Anderson. G.J. Transfusion suppresses erythropoiesis and increases hepcidin in adult patients with beta-thalassemia major: a longitudinal study. *Blood*. **2013**, *122*, *1*, 124-133.
23. Girard, J.M., Drevin, G., Brasme, J.F., Pissard, S., Reynier, P., Simard, G. and Denis, M.C., 2016, November. Clinical and biological specificity of beta-thalassemia intermedia: a case report. *Ann Biol Clin (Paris)*.**2016**,*74*, *6*, 688-692.
24. Jamieson. C.; Hasserjian. R.; Gotlib. J.; Cortes. J.; Stone. R.; Talpaz. M.; Thiele. J.; Rodig. S.; Pozdnyakova. O. Effect of treatment with a JAK2-selective inhibitor, fedratinib, on bone marrow fibrosis in patients with myelofibrosis. *J Transl Med*. **2015**, *13*, *1*, 294-301.
25. Taher, A.; Vichinsky, E.; Musallam, K.; Cappellini. M.D.; Viprakasit, V. Guidelines for the management of non-transfusion dependent thalassaemia (NTDT). Thalassaemia International Federation, Nicosia, Cyprus; **2013**.
26. Al-Ali. H.K.; Stalbovskaya. V.; Gopalakrishna.P.; Perez-Ronco. J.; Foltz. L. Impact of ruxolitinib treatment on the hemoglobin dynamics and the negative prognosis of anemia in patients with myelofibrosis. *Leuk. Lymphoma*. **2016**, *57*, *10*, 2464–2467.
27. Zhao. N.; Nizzi. C.P.; Anderson. S.A.; Wang. J.; Ueno. A.; Tsukamoto. H.; Eisenstein. R.S.; Enns. C.A.; Zhang. A.S. Low intracellular iron increases the stability of matriptase-2. *J. Biol. Chem*. **2015**, *290*, *7*, 4432–4446.
28. Wu. X.G.; Wang.Y.; Wu. Q.; Cheng. W.H.; Liu. W.; Zhao. Y.; Mayeur. C.; Schmidt. P.J.; Paul. B.Y.; Wang. F.; Xia. Y. HFE interacts with the BMP type I receptor ALK3 to regulate hepcidin expression. *Blood*. **2014**, *124*, 1335–1343.
29. Rishi. G.; Crampton. E.M.; Wallace. D.F.; Subramaniam. V.N. In situ proximity ligation assays indicate that hemochromatosis proteins Hfe and transferrin receptor 2 (Tfr2) do not interact. *PLoS One*. **2013**, *8*, *10*, 1-11.