



## New Travelling Wave Solution of *Burgers Equations*

**Zainab John**

Department of Mathematical, College of  
Science, Al-Mustansiriyah University,  
Baghdad, Iraq.

[zainabjohn22@uomustansiriyah.edu.iq](mailto:zainabjohn22@uomustansiriyah.edu.iq)

**Basim Akhudir Abbas**

Department of Mathematical, College of  
Science, Al-Mustansiriyah University,  
Baghdad, Iraq.

[baasim\\_math@uomustansiriyah.edu.iq](mailto:baasim_math@uomustansiriyah.edu.iq)

**Article history: Received 19 January 2021, Accepted 11 April 2021, Published in 2021.**

**Doi: 10.30526/34.3.2678**

### Abstract

In this paper, we studied the travelling wave solving for some models of Burger's equations. We used sine-cosine method to solution nonlinear equation and we used a direct solution after getting travelling wave equation.

**Keywords:** travelling wave solution, cosine-sine Method, nonlinear differential equations, Wave equations.

### 1.Introduction

A large variety of physical, chemical, and biological phenomena are waves, such as sound waves, string waves, water waves, etc. Wave phenomena are very important in dispersion, diffusion, reaction, and convection, and such phenomena are represented by nonlinear partial differential equations.

Moving waves are visible in many linear equations and nonlinear wave modeling, such as heat waves, string waves, and the cycle life of some organisms one of the travelling waves, etc. One of the most important methods for solving wave equations is the Travelling Wave solution (TWS), which many researchers have discussed ( see [1,2], the tanh-coth method [3], the tanh [4,5] sine-cosine method [6], and see [7-12]).

### 2.Infinite Sires Method [5,9]

To clarify the travelling wave solutions method, we note the following:

Let the nonlinear partial differential equation has the form

$$g(u, u_t, u_x, u_{xx}, \dots) = 0, \quad (1)$$



Where  $u = (x, t)$  is a solution of equation (1), and  $g$  is a polynomial with respect to  $u$  and with its derivatives. To solve Eq. (1), we follow the following steps:

**Step 1:** First, we change the independent variables  $x, t$ , by using one independent variable  $\xi$  that combines the two variables as shown

$$\text{Let } u(x, t) = f(\xi) \quad \text{where } \xi = x - ct, \quad (2)$$

Where  $\xi$  is a wave variable and  $c$  is constant.

**Step2:** derivative equation (2) with respect to  $x$  and  $t$ , we have the following ordinary differential equations:

$$u_x = u' = \frac{du}{d\xi}; u_{xx} = u'' = \frac{d^2u}{d\xi^2}; u_t = -cu' = -c \frac{du}{d\xi}; u_{tt} = c^2u'' = c^2 \frac{d^2u}{d\xi^2};$$

$$u_{xt} = -cu'' = -c \frac{d^2u}{d\xi^2} \quad (3)$$

When substitute (3) into (1), we find that equation (1) is transform to (linear or nonlinear) ordinary differential equation transformed into

$$G\left(f(\xi), \frac{df}{d\xi}, \frac{d^2f}{d\xi^2}, \dots\right) \quad (4)$$

Where  $G$  is a polynomial in  $f(\xi)$  and with its derivatives,

**Step 3:** integrate (4) with respect to  $\xi$ , and let the constant of integration equal to zero.

**Step 4: (a)** we can solve equation (4) directly through many methods that solve ordinary differential equations.

or

**(b)** use sine-cosine method to solve equation (4).

### 3. Applications of the travelling wave solution

#### 3.1: Some Models of Burgers Equations [13]

The viscid Burgers equation to be the nonlinear parabolic PDE has the form

$$v_t = v_{xx} + \epsilon vv_x, \quad 0 < \epsilon \leq 1. \quad (5)$$

##### 3.1.1. Travelling wave solution

The travelling wave solution of Eq.(5) is

$$\xi = x - ct \Rightarrow v(x, t) = v(\xi), \quad v_x = v' = \frac{dv}{d\xi}; v_{xx} = v'' = \frac{d^2v}{d\xi^2}; v_t = -cv' = -c \frac{dv}{d\xi}. \quad (6)$$

Substituting Eq. (6) into Eq. (5), we have got

$$v'' + cv' + \epsilon vv' = 0 \quad (7)$$

$$v'' + cv' + \epsilon \left(\frac{v^2}{2}\right)' = 0 \quad (8)$$

Integral Eq. (8) with respect to  $\xi$  we have got

$$v' + cv + \epsilon \frac{v^2}{2} = c_1 \quad (9)$$

Let  $c_1 = 0$

$$v' = -(cv + \epsilon \frac{v^2}{2}) \tag{10}$$


$$\frac{dv}{-(cv + \epsilon \frac{v^2}{2})} = d\xi$$

By solving this equation, we have got

$$v(\xi) = \frac{c}{(ce^{c(\xi+c_2)} - \epsilon/2)} \tag{11}$$

Where  $\xi = x - ct$ , substituted in (11), we have  $v(x, t) = \frac{c}{(ce^{(x-ct)+c_2} - \epsilon/2)}$ , which is a general solution of Burgers equation (5).

By drawing the solution function  $(v(x,t))$ , we find the following figure



Singular kink of u

Figure 1. When  $c=1, \delta=1/3, x=[-10,10], t=[-10,10]$ .

### 3.2. KdV-Burgers Equation:

#### 3.2.1: Cosine Function Method:

In this part, we want to find (TWS) of KdV-Burgers equation, by using the cosine function method see [6,7]. KdV-Burgers equation [13,14] has the form

$$u_t + (u^2)_x - \gamma u_{xx} + \delta u_{xxx} = 0, \tag{12}$$

Where  $\delta$  and  $\gamma > 0$  are constants and  $u$  is a function of spatial variable  $x$  and time variable  $t$

$$\text{let } u(x, t) = u(\xi), \quad \xi = x - ct \tag{13}$$

Application (TWS) on Eq. (12) yields into the ordinary differential equation

$$-cu' + (u^2)' - \gamma u'' + \delta u''' = 0 \tag{14}$$

Integrating (14) we get

$$-cu + u^2 - \gamma u' + \delta u'' = 0 \tag{15}$$

Then we have the nonlinear travelling wave equation (15). Solution Eq.+(15) by cosine function method has the form.

$$u(\xi) = \lambda \cos^\beta(\mu\xi) \tag{16}$$

Where  $\lambda$ ,  $\beta$  and  $\mu$  are unknown parameters to find its, we derivatives (16) we get

$$u' = -\lambda\mu\beta\cos^{\beta-1}(\mu\xi)\sin(\mu\xi) \tag{17}$$

$$u'' = -\lambda\beta\mu^2\cos^\beta(\mu\xi) + \lambda\mu^2\beta(\beta - 1)\cos^{\beta-2}(\mu\xi). \tag{18}$$

Substituting (16-18) into equation (15) gives

$$-c\lambda\cos^\beta(\mu\xi) + (\lambda\cos^\beta(\mu\xi))^2 + \gamma\lambda\mu\beta\cos^{\beta-1}(\mu\xi)\sin(\mu\xi) + \delta(-\lambda\beta\mu^2\cos^\beta(\mu\xi) + \lambda\mu^2\beta(\beta - 1)\cos^{\beta-2}(\mu\xi)) = 0$$

The following algebraic equation system is obtained by equating the exponents and the coefficient of each pair of the cosine functions.

$$-c\lambda - \delta\lambda\beta\mu^2 = 0 \tag{19}$$

$$\lambda^2 + \delta\lambda\mu^2\beta(\beta - 1) = 0 \tag{20}$$


$2\beta = \beta - 2 \Rightarrow \beta = -2$ , substitute in (19-20) we have

$$\mu = \mp \sqrt{\frac{c}{2\delta}}, \quad \lambda = 3c, \text{ substitute in (16) we get the solution}$$

$$u(\xi) = 3c \cos^{-2}\left(\sqrt{\frac{c}{2\delta}} \xi\right) \text{ or } u(\xi) = -3c \cos^{-2}\left(\sqrt{\frac{c}{2\delta}} \xi\right)$$

$$u(x,t) = 3c \sec^2\left(\sqrt{\frac{c}{2\delta}}(x - ct)\right) \tag{21}$$

By drawing the solution function  $u(x,t)$ , we find the following figure



The periodic solution of u  
**Figure 2.** When  $c=1$ ,  $\delta=1/3$ ,  $x=[-10,10]$ ,  $t=[-10,10]$ .

**3.2.2.Sine Function Method:**

Solve equation (15) by use (sine-function method), we have the form,

$$u = \lambda \sin^\beta(\mu\xi) \tag{22}$$

Where  $\lambda$ ,  $\beta$  and  $\mu$  are unknown parameters, to find its, we Derivatives (22) we get.

$$u' = \lambda\mu\beta \sin^{\beta-1}(\mu\xi)\cos(\mu\xi) \tag{23}$$

$$u'' = -\lambda\beta\mu^2 \sin^\beta(\mu\xi) + \lambda\mu^2\beta(\beta - 1)\sin^{\beta-2}(\mu\xi) \tag{24}$$

Substituting (22-24) into equation (15) yield in

$$-c\lambda \sin^\beta(\mu\xi) + \left(\lambda \sin^\beta(\mu\xi)\right)^2 - \gamma\lambda\mu\beta \sin^{\beta-1}(\mu\xi)\cos(\mu\xi) + \delta(-\lambda\beta\mu^2 \sin^\beta(\mu\xi) + \lambda\mu^2\beta(\beta - 1)\sin^{\beta-2}(\mu\xi)) = 0$$

We find the following system of algebraic and we obtain to parameters by equating the exponents and the coefficient of each pair of the (cosine function).

$$-c\lambda - \delta\lambda\beta\mu^2 = 0 \tag{25}$$

$$\lambda^2 + \lambda\mu^2\beta(\beta - 1) = 0, \tag{26}$$

$$2\beta = \beta - 2 \Rightarrow \beta = -2 \tag{27}$$


Substitute (27) in (25,26) we have

$$\mu = \mp \sqrt{\frac{c}{2\delta}}, \lambda = 3c$$

Then, we have the solution

$$u = -3c \sin^{-2}\left(\sqrt{\frac{c}{2\delta}}\xi\right) \text{ or } u = 3c \sin^{-2}\left(\sqrt{\frac{c}{2\delta}}\xi\right)$$

$$u = \mp 3c \csc^2\left(\sqrt{\frac{c}{2\delta}}(x - ct)\right)$$



The periodic solutions of u  
**Figure 3.** When  $c=1$ ,  $\delta=1/3$ ,  $x=[-10,10]$ ,  $t=[-10,10]$ .

### Conclusion

In this work, we conclude that the method of sine and cosine function gives the exact solution to the nonlinear partial differential equation, which is difficult to find in other ways.

### References

1. Hirota, R. Exact envelope-soliton solutions of a nonlinear wave equation. *Journal of Mathematical Physics*, **1973**, 14(7), 805-809.
2. Hirota, R., ; Satsuma, J. Soliton solutions of a coupled Korteweg-de Vries equation. *Physics Letters A*, **1981**, 85(8-9), 407-408.
3. Kumar, A. ; Pankaj, R. D. Tanh-coth scheme for traveling wave solutions for Nonlinear Wave Interaction model. *Journal of the Egyptian Mathematical Society*, 23(2), 282-285.
4. Taha, W. M., Hameed, R. A. ; Noorani, M. S. M. (2016). Traveling Wave Solutions by using Extended Tanh Function Method with special values. *Diyala Journal for Pure Science*, **2015**, 12 (4-part 2), 1-13.
5. Fan, E. Extended tanh-function method and its applications to nonlinear equations. *Physics Letters A*, **2000**, 277(4-5), 212-218.
6. Bibi, S., ; Mohyud-Din, S. T. Traveling wave solutions of KdVs using sine-cosine method. *Journal of the Association of Arab Universities for Basic and Applied Sciences*, **2014**, 15(1), 90-93.
7. Taghizadeh, N., Mirzazadeh, M. ; Farahrooz, F. Exact travelling wave solutions of the coupled Klein-Gordon equation by the infinite series method. *Applications and Applied Mathematics: An International Journal (AAM)*, **2011**, 6, 1964-1972.

8. He, J. H. ; Wu, X. H. Exp-function method for nonlinear wave equations. *Chaos, Solitons & Fractals*. **2006**, 30(3), 700-708.
9. Bekir, A., & Boz, A. Exact solutions for nonlinear evolution equations using Exp-function method. *Physics Letters A*, **2008**, 372(10), 1619-1625.
10. Alam, M. N., & Akbar, M. A. Traveling wave solutions for the mKdV equation and the Gardner equations by new approach of the generalized (G'/G)-expansion method. *Journal of the Egyptian Mathematical Society*, **2014**, 22(3), 402-406.
11. Bibi, S., & Mohyud-Din, S. T. New traveling wave solutions of Drinefel'd–Sokolov–Wilson Equation using Tanh and Extended Tanh methods. *Journal of the Egyptian Mathematical Society*, **2014**, 22(3), 517-523.
12. Shi, Q., Xiao, Q., & Liu, X. Extended wave solutions for a nonlinear Klein–Gordon–Zakharov system. *Applied Mathematics and Computation*, (2012). 218(19), 9922-9929.
13. Wazwaz. A.M. Partial Differential Equations: Method and Applications, *Taylor and Francis*, London. **2002**.
14. Wang, M. L. Exact solution for a compound KdV-Burgers equation. *Phys*,(1966). *Lett. A* 213(279-287).