



## Some Properties of Connectedness in Grill Topological Spaces

**Saad S. Suliman**

Department of Mathematics, College of  
Education for Pure Science, Ibn Al  
Haitham, University of Baghdad, Iraq.  
[saadsadeq05@gmail.com](mailto:saadsadeq05@gmail.com)

**Esmaeel R.B.**

Department of Mathematics, College  
of Education for Pure Science, Ibn Al  
Haitham, University of Baghdad, Iraq.  
[Ranamumosa@yahoo.com](mailto:Ranamumosa@yahoo.com)

**Article history: Received 12 June 2022, Accepted 21 August 2022, Published in October 2022.**

**Doi: 10.30526/35.4.2878**

### Abstract

We use the idea of Grill, this study generalized a new sort of linked space like  $\alpha$ -connected – hyperconnected and investigated its features, as well as the relationship between it and previously described notation. It also developed new sorts of functions, such as hyperconnected space, and identifying their relationship, by offering numerous instance and attributes that belong to this set. This set will serve as a starting point for further research into the sets many future possibilities. Also, we use some of the theorems and observations previously studied and relate them to the grill and the Alpha group, and benefit from them in order to obtain new results in this research. We applied the concept of Connected to them and obtained results related to Connected. The sources related to the Connected and Alpha were considered as starting points and an important basis in this research.

**Keywords:** Grill,  $\mathbb{G}^* \alpha$ -connected,  $\mathbb{G}^* \alpha$ -disconnected,  $\mathbb{G}^* \alpha o(\mathbb{W})$ ,  $\mathbb{G}^* \alpha$ -open.

### 1. Introduction

[1], [2] established a grill notion in a topological space, and the grill has shown to be an effective tool for learning a variety of topological concerns. A subset of a topological space  $(\mathbb{W}, \tau)$  which is a non-empty collection  $\mathbb{G}$  is referred to be a grill whenever (a)  $\mathbb{A} \in \mathbb{G}$  and  $\mathbb{A} \subseteq \mathbb{M}$  implying  $\mathbb{B} \in \mathbb{G}$ . (b)  $\mathbb{W}$  has a subset  $\mathbb{A}$  and  $\mathbb{M}$  also  $\mathbb{A} \cup \mathbb{B} \in \mathbb{G}$  lead to  $\mathbb{A} \in \mathbb{G}$  or  $\mathbb{B} \in \mathbb{G}$ . A triple  $(\mathbb{W}, \tau, \mathbb{G})$  topological space with grills is one type of topological space.

Mukherjee and Roy [3] created a distinctive topology with a grill and investigated topological notions. For every topological space  $(\mathbb{W}, \tau)$  point  $\mathbb{W}$ , Neighborhoods are open of  $\mathbb{W}$  embodied



by  $\tau(\mathbb{W})$ . A mapping  $\Psi: \mathbb{P}(\mathbb{W}) \rightarrow \mathbb{P}(\mathbb{W})$  is referred to as  $\phi(\mathbb{A}) = \{ \mathbb{W} \in \mathbb{W}: \mathbb{A} \cap \tilde{\mathbb{U}} \in \mathbb{G} \text{ apiece } \tilde{\mathbb{U}} \in \tau(\mathbb{W}) \}$  for every  $\mathbb{A} \in \mathbb{P}(\mathbb{W})$ . A mapping  $\Psi: \mathbb{P}(\mathbb{W}) \rightarrow \mathbb{P}(\mathbb{W})$  is referred to as  $\Psi(\mathbb{A}) = \mathbb{A} \cup \phi(\mathbb{A})$  for every  $\mathbb{A} \in \mathbb{P}(\mathbb{W})$ . The map  $\Psi$  Kuratowski closure axioms are met:

- (a)  $\Psi(\emptyset) = \emptyset$ ,
- (b) when  $\mathbb{A} \subseteq \mathbb{B}$ , so  $\Psi(\mathbb{A}) \subseteq \Psi(\mathbb{B})$ ,
- (c) when  $\mathbb{A} \subseteq \mathbb{W}$ , so  $\Psi(\Psi(\mathbb{A})) = \Psi(\mathbb{A})$ ,
- (d) when  $\mathbb{A}, \mathbb{B} \subseteq \mathbb{W}$ , so  $\Psi(\mathbb{A} \cup \mathbb{B}) = \Psi(\mathbb{A}) \cup \Psi(\mathbb{B})$ .

Grill topological spaces come in a variety of shapes and size as a discrete topology and a complement finite topology.

[4] [5] On a space  $(\mathbb{W}, \tau)$ , this agrees to a grill  $\mathbb{G}$  a topology exists  $\tau_{\mathbb{G}}$  on  $\mathbb{W}$  that is given by no one else by  $\tau_{\mathbb{G}} = \{ \tilde{\mathbb{U}} \subseteq \mathbb{W}: \Psi(\mathbb{W} - \tilde{\mathbb{U}}) = \mathbb{W} - \tilde{\mathbb{U}} \}$ , Consequently,  $\mathbb{A} \subseteq \mathbb{W}$ ,  $\Psi(\mathbb{A}) = \mathbb{A} \cup \phi(\mathbb{A})$ .  $\tau \subseteq \tau_{\mathbb{G}}$  and  $\Psi(\mathbb{A}) = cl(\mathbb{A})$ . Using the following as a basis, we are able to locate  $\tau_{\mathbb{G}}$  on  $\mathbb{W}$  by supplying  $\tau_{\mathbb{G}}$  through using the following as a basis  $\beta(\tau_{\mathbb{G}}, \mathbb{W}) = \{ \mathbb{Y} - \mathbb{A}; \mathbb{Y} \in \tau, \mathbb{A} \notin \mathbb{G} \}$ .

[6]. On a space  $(\mathbb{W}, \tau)$ , there is a grill  $\mathbb{G}$ ,  $\tau \subseteq \beta(\mathbb{G}, \tau) \subseteq \tau_{\mathbb{G}}$ , where  $\beta(\mathbb{G}, \tau)$  basis for  $\tau_{\mathbb{G}}$ . As an example, [7] to exist in a space  $(\mathbb{W}, \tau)$ ,  $\tau_{\mathbb{G}} = \tau$  whenever  $\mathbb{G} = \mathbb{P}(\mathbb{W}) \setminus \{ \emptyset \}$  implying  $\tau_{\mathbb{G}} = \tau$ . The family of all  $\alpha$ -open set is showed by  $\tau_s$ .

$\alpha$ -open is a subset  $\mathbb{A}$  of a space  $(\mathbb{W}, \tau)$ , [8] if  $\mathbb{A} \subseteq cl(int(\mathbb{A}))$ ,

Let  $(\mathbb{W}, \tau, \mathbb{G})$  be topological space. The subset  $\mathbb{A}$  in  $\mathbb{W}$  is known as  $\mathbb{G}$   $\alpha$ -open if  $\mathbb{A} \subseteq \Psi(int(\mathbb{A}))$ , and every  $\Psi \alpha$ -open is an  $\alpha$ -open. Many researchers have generalized using these [9] [10]. The symbol  $int(\mathbb{A})$  is the interior set of  $\mathbb{A}$  and  $cl(\mathbb{A})$  denotes the closure of  $\mathbb{A}$ .

combinations

The space  $(\mathbb{W}, \tau)$  is disconnected if and only if there exist two open disjoint nonempty sets  $\mathbb{A}$  and  $\mathbb{B}$  such that  $\mathbb{A} \cup \mathbb{B} = \mathbb{W}$ . i.e.,  $\mathbb{W}$  is disconnected if and only if  $\mathbb{W} = \mathbb{A} \cup \mathbb{B}; \mathbb{A}, \mathbb{B} \in \tau, \mathbb{A} \cap \mathbb{B} = \emptyset, \mathbb{A}, \mathbb{B} \neq \emptyset$ .

The sets  $\mathbb{A}$  and  $\mathbb{B}$  form a separation of  $\mathbb{W}$ . The space  $(\mathbb{W}, \tau)$  is connected if and only if it is not disconnectaed.  $\mathbb{W}$  is connected if and only if  $\mathbb{W} \neq \mathbb{A} \cup \mathbb{B}; \mathbb{A}, \mathbb{B} \in \tau, \mathbb{A} \cap \mathbb{B} = \emptyset, \mathbb{A}, \mathbb{B} \neq \emptyset$ .

## 2. Grill $\alpha$ -open sets

### Definition 2.1.

In [10], the set  $\mathbb{A}$  is referred to as Grill  $\alpha$ -open if  $\mathbb{A} \in \tau; \mathbb{A} - \Psi(int(\mathbb{A})) \notin \mathbb{G}$  and  $\mathbb{A} - int(cl(\mathbb{A})) \notin \mathbb{G}$ . As stated by  $\mathbb{G}^* \alpha$ -open, the complment of  $\mathbb{G}^* \alpha$ -open is  $\mathbb{G}^* \alpha$ -closed. The set of all  $\mathbb{G}^* \alpha$ -open is represented by  $\mathbb{G}^* \alpha o(\mathbb{W})$ , and  $\mathbb{G}^* \alpha$ -closed is denoted by  $\mathbb{G}^* \alpha c(\mathbb{W})$ .

### Example 2.2. [10]

Let  $(\mathbb{W}, \tau, \mathbb{G})$  is a topological space of the grill, and let  $\mathbb{W} = \{ w_1, w_2, w_3 \}, \tau = \{ \mathbb{W}, \emptyset, \{ w_1 \}, \{ w_1, w_2 \} \}, \mathcal{F} = \{ \mathbb{W}, \emptyset, \{ w_3 \}, \{ w_2, w_3 \} \}, \mathbb{G} = \{ \mathbb{S} \subseteq \mathbb{W}; w_2 \in \mathbb{S} \}, \emptyset: \mathbb{P}(\mathbb{W}) \rightarrow \mathbb{P}(\mathbb{W}), \emptyset(\mathbb{A}) = \{ \mathbb{W} \in \mathbb{W}; \forall \mathbb{S} \in \tau_x; \mathbb{S} \cap \mathbb{A} \in \mathbb{G} \}, \Psi(\mathbb{A}) = \mathbb{A} \cup \emptyset, \tau_{\mathbb{G}} = \{ \mathbb{W}, \emptyset, \{ w_2, w_1 \}, \{ w_3, w_2 \}, \{ w_2 \} \}, \mathcal{F}_{\mathbb{G}} = \{ \mathbb{W}, \emptyset, \{ w_3 \}, \{ w_1 \}, \{ w_2, w_3 \} \}, so  $\mathbb{G}^* \alpha o(\mathbb{W}) = \{ \mathbb{W}, \emptyset, \{ w_3 \}, \{ w_2 \}, \{ w_1 \}, \{ w_2, w_3 \}, \{ w_1, w_2 \}, \{ w_3, w_1 \} \}$ .$

**Theorem 2.3.** [10]

The union of any family of  $\mathbb{G}^* - \alpha$  open set is a  $\mathbb{G}^* - \alpha$  open set.

**Remark 2.4.** [10]

Both ideas are related  $\mathbb{G}^* \alpha -$  open set ,  $\alpha -$  open set are independent.

**Remark 2.5.** [10]

Supra topology refers to the collection of all  $\mathbb{G}^* \alpha$ -open sets.

**Remark 2.6** [10]

- i. The open set leads to  $\mathbb{G}^* \alpha$ - open set.
- ii. The closed set leads to  $\mathbb{G}^* \alpha$ - closed set.

**Theorem 2.7.** [10]

The  $\mathbb{G} \alpha$ -open leads to  $\mathbb{G}^* \alpha$ -open.

**Proposition 2.8.** [10]

For any Grill topology  $(\mathbb{W}, \tau, \mathbb{G})$ ,  $\mathbb{A}$  is a  $\mathbb{G} \alpha$ -open set if and only if  $\mathbb{A}$  is a  $\mathbb{G}^* \alpha$ -open set whenever  $\mathbb{G} = \mathbb{P}(\mathbb{W}) \setminus \{\emptyset\}$ .

**Definition 2.9.** [10]

The function  $f: (\mathbb{W}, \tau, \mathbb{G}) \rightarrow (\mathbb{Y}, \tau', \mathbb{G})$  is referred to as:

- 1.  $\mathbb{G}^* \alpha$ -open function, shortly " $\mathbb{G}^* \alpha$ -o function" if  $f(\mathbb{s}) \in \mathbb{G}^* \alpha O(\mathbb{y})$  when  $\mathbb{s} \in \mathbb{G}^* \alpha O(\mathbb{W})$ .
- 2.  $\mathbb{G}^{**} \alpha$ - open function, shortly " $\mathbb{G}^{**} \alpha$ -o function " if  $f(\mathbb{s}) \in \mathbb{G}^* \alpha O(\mathbb{y})$  when  $\mathbb{s} \in \tau$ .
- 3.  $\mathbb{G}^{***} \alpha$ - open function, shortly " $\mathbb{G}^{***} \alpha$ -o function " if  $f(\mathbb{s}) \in \tau'$  whenever  $\mathbb{s} \in \mathbb{G}^* \alpha O(\mathbb{W})$ .

**Definition 2.10**

The function  $f: (X, \tau, \mathbb{G}) \rightarrow (\mathbb{W}, \tau', \mathbb{G})$  is called ;

- 1.  $\mathbb{G}^* \alpha -$ continuouse function, shortly " $\mathbb{G}^* \alpha$ -continuous function" " if  $f^{-1}(\mathbb{s}) \in \mathbb{G}^* \alpha O(X)$  for all  $\mathbb{s} \in \tau'$ .
- 2. Strongly  $\mathbb{G}^* \alpha -$  continuouse function shortly "strongly  $\mathbb{G}^* \alpha$ -continuous function" if  $f^{-1}(\mathbb{s}) \in \tau$ , fore every  $\mathbb{s} \in \mathbb{G}^* \alpha O(\mathbb{W})$ .
- 3.  $\mathbb{G}^* \alpha -$ irresolute function, shortly " $\mathbb{G}^* \alpha$ -irresolute function" if  $f^{-1}(\mathbb{s}) \in \mathbb{G}^* \alpha O(X)$ , for every  $\mathbb{s} \in \mathbb{G}^* \alpha O(\mathbb{W})$

**3. Grill  $\alpha$  -open sets in grill connected space**

**Definition 3.1:**

The space  $(\mathbb{W}, \tau, \mathbb{G})$  is a  $\mathbb{G}^* \alpha O$ - disconnected if and only if there exist two  $\mathbb{G}^* \alpha$ -open disjoint nonempty sets  $\mathbb{A}$  and  $\mathbb{B}$  such that  $\mathbb{A} \cup \mathbb{B} = \mathbb{W}$ . i.e.,  $\mathbb{W}$  is a  $\mathbb{G}^* \alpha O$ - disconnected if and only if  $\mathbb{W} = \mathbb{A} \cup \mathbb{B}$  ;  $\mathbb{A}, \mathbb{B} \in \mathbb{G}^* so(\mathbb{W})$  ,  $\mathbb{A} \cap \mathbb{B} = \phi$ .

The sets  $\mathbb{A}$  and  $\mathbb{B}$  form a  $\mathbb{G}^* so$ -separation of  $\mathbb{W}$ . The space  $(\mathbb{W}, \tau, \mathbb{G})$  is a  $\mathbb{G}^* so$ - connected if and

only if it is not  $\mathbb{C}^* \alpha o$ - disconnected.  $W$  is a  $\mathbb{C}^* \alpha o$ - connected if and only if  $W \neq A \cup B$  ;  $A, B \in \mathbb{C}^* \alpha o(W)$ ,  $A \cap B = \phi$ ,  $A, B \neq \phi$ .

**Example 3.2.**

Let  $(W, \tau, \mathbb{C})$  topological space be a grill and  $W = \{W_1, W_2, W_3\}$ ,  $\tau = \tau_{\mathbb{C}} = \{W, \emptyset, \{W_2\}, \{W_3\}\}$ ,  $\mathbb{C} = \mathbb{P}(W) \setminus \{\emptyset\}$  is a  $\mathbb{C}^* \alpha o$ - connected space.

**Remark 3.3**

Every disconnected set is a  $\mathbb{C}^* \alpha o$ -disconnected.

**Proof.**

Let  $W$  is disconnected set, then the exist  $W \neq \emptyset, Z \neq \emptyset$  and  $W, Z \in \tau$ ;  $W \cap Z = \emptyset$  and  $W \cup Z = W$  since every open set in  $\mathbb{C}^* \alpha$ -open set. Therefore,  $W$  is  $\mathbb{C}^* \alpha o$ -disconnected.

**Remark 3.4.**

The space  $(W, \tau, \mathbb{C})$  is a  $\mathbb{C}^* \alpha o$ - disconnected with any grill and  $\mathbb{C}^* \alpha o(W) = \mathbb{P}(W)$  if  $W$  contained more than one element since there exist  $A$  and  $A^c \in \mathbb{C}^* \alpha o(W)$ ;  $A \cup B = W$ ,  $A \cap B = \phi$ ,  $A, B \neq \phi$ .

**Remark 3.5.**

The space  $(W, \tau, \mathbb{C})$  is a  $\mathbb{C}^* \alpha o$ - disconnected and  $\mathbb{C} = \emptyset$  and  $\tau_{\mathbb{C}} = \{W, \emptyset\}$ , so  $\mathbb{C}^* \alpha o$ - connected space.

**Remark 3.6.**

If  $\tau_{\mathbb{C}} = \mathcal{F}_{\mathbb{C}}$  and  $\tau_{\mathbb{C}} \neq \text{Indiscret}$  , when  $\mathbb{C} = \mathbb{P}(W) \setminus \{\emptyset\}$ . This means that  $(W, \tau, \mathbb{C})$  is a  $\mathbb{C}^* \alpha o$ - disconnected .

**Theorem 3.7.**

The space  $(W, \tau, \mathbb{C})$  is a  $\mathbb{C}^* \alpha o$ -connected if and only  $W$  cannot be written as a union of two non-empty disjoint closed sets.

**Proof.**

Let  $W$  be a  $\mathbb{C}^* \alpha o$ -connected if  $W = A \cup B$ , such that;  $A$  and  $B \in \mathbb{C}^* \alpha c(W)$  ,  $A \cap B = \phi$ ,  $A, B \neq \phi$  , so.  $A = B^c$  and  $B = A^c$  , then  $W = A \cup B$ ;  $A$  and  $B \in \mathbb{C}^* \alpha o(W)$ ,  $A \cap B = \phi$ ,  $A, B \neq \phi$  , that is mean that  $W$  is a  $\mathbb{C}^* \alpha o$ - disconnected. That is contradiction. Therefore,  $W$  cannot be written as a union of two non-empty disjoint closed set. Now, if  $W$  is a  $\mathbb{C}^* \alpha o$ - disconnected space ,so.  $W = A \cup B$ ;  $A$  and  $B \in \mathbb{C}^* \alpha o(W)$ ,  $A \cap B = \phi$ ,  $A, B \neq \phi$  , so  $A = B^c$  and  $B = A^c$ , then  $A$  and  $B \in \mathbb{C}^* \alpha c(x)$ , but that is a contradiction. Therefore,  $W$  is a  $\mathbb{C}^* \alpha o$ - connected space.

**Theorem 3.8.**

The space  $(W, \tau, \mathbb{C})$  is a  $\mathbb{C}^* \alpha o$ -connected if and only if the only the subsets of the space  $W$  which are  $\mathbb{C}^* \alpha$ -open and  $\mathbb{C}^* \alpha$ -closed are  $W$  and  $\emptyset$  .

**Proof.**

Let  $A$  and  $A^c \in \mathbb{C}^* \alpha o(x)$  and  $A \neq Y, A \neq \emptyset$ , so  $Y = A \cup A^c; A \cap A^c = \phi, A, A^c \neq \phi$ . Then,  $Y$  is a  $\mathbb{C}^* \alpha o$ -disconnected, and that is a contradiction. So, where  $A \subseteq Y, A, A^c \in \mathbb{C}^* \alpha o(x)$ , then  $A = Y$  or  $A = \emptyset$ , let  $Y$  be a  $\mathbb{C}^* \alpha o$ -disconnected space. This means that  $Y = W \cup Z; W, Z \in \mathbb{C}^* \alpha o(x), W \cap Z = \phi$  and  $W, Z \neq \emptyset$  implies that  $W = Z^c$  and  $Z = W^c$ . So  $W, Z \in \mathbb{C}^* \alpha c(x)$  (that is contradiction) Therefore,  $Y$  is a  $\mathbb{C}^* \alpha o$ -connected space.

**Remark 3.9.**

If  $f: (Y, \tau, \mathbb{C}) \rightarrow (Y, \tau', \mathbb{C})$  is a  $\mathbb{C}^* \alpha o$ -irresolute and onto function and  $Y$  is a  $\mathbb{C}^* \alpha o$ -connected space then  $Y$  is not necessary  $\mathbb{C}^* \alpha o$ -connected space.

**Example 3.10.**

Let  $f: (\mathbb{R}, D, \mathbb{C}) \rightarrow (\mathbb{R}, I, \mathbb{C}); f(W) = W$  for all  $W \in \mathbb{R}$  and  $\mathbb{C} = \mathbb{P}(W) \setminus \{\emptyset\}$ , so  $f$  is a  $\mathbb{C}^* \alpha o$ -irresolute and onto function and  $(\mathbb{R}, I, \mathbb{C})$  is a  $\mathbb{C}^* \alpha o$ -connected space and  $(\mathbb{R}, D, \mathbb{C})$  is a  $\mathbb{C}^* \alpha o$ -disconnected space.

**Theorem 3.11.**

If  $A$  and  $B$  are a  $\mathbb{C}^* \alpha o$ -connected spaces of  $(Y, \tau, \mathbb{C})$  and  $A \cap B \neq \phi$ , then  $A \cup B$  is a  $\mathbb{C}^* \alpha o$ -connected space.

**Proof.**

Let  $(Y, \tau, \mathbb{C})$  be a grill topological space and  $A, B \subseteq Y; A, B$  be a  $\mathbb{C}^* \alpha o$ -connected space. Now, If  $A \cup B$  is a  $\mathbb{C}^* \alpha o$ -disconnected space, so  $A \cup B = W \cup Z; W, Z \in \mathbb{C}^* \alpha o(x)_{(A \cup B)}, W \cap Z = \phi$  and  $W, Z \neq \emptyset$ , then  $A \subseteq A \cup B, A \subseteq W \cup Z, A \subseteq W$  or  $A \subseteq Z$ . Similarly,  $B$  leads to either  $A \subseteq W$  and  $B \subseteq W$ , then  $A \cup B \subseteq W$ , then  $Z = \emptyset$  contradiction. Or,  $A \subseteq Z$  and  $B \subseteq Z$ , then  $A \cup B \subseteq Z$ , then  $W = \emptyset$  C! Or  $A \subseteq W$  and  $B \subseteq Z$ , then  $A \cap B \subseteq W \cap Z$ , then  $A \cap B = \emptyset$  C! Or  $A \subseteq Z$  and  $B \subseteq W$ , then  $A \cap B \subseteq W \cap Z$ , then  $A \cap B = \emptyset$ . This is a contradiction. So,  $A \cup B$  is a  $\mathbb{C}^* \alpha o$ -connected space.

**Remark 3.12.**

We can generalize **Theorem 3.11** to a family of a  $\mathbb{C}^* \alpha o$ -connected sets as follows: let  $\{A_\alpha\}_{\alpha \in \Lambda}$  be a family of a  $\mathbb{C}^* \alpha o$ -connected subsets of a space  $(Y, \tau, \mathbb{C})$  and  $\bigcap_{\alpha \in \Lambda} A_\alpha \neq \emptyset$ , then  $\bigcup_{\alpha \in \Lambda} A_\alpha$  is a  $\mathbb{C}^* \alpha o$ -connected set.

**4. Grill  $\alpha$ -open sets in grill connected space hyperconnected**

**Definition 4.1.**

In any grill topological space,  $(Y, \tau, \mathbb{C})$  is said to be:

1.  $*$ -hyperconnected if  $A$  is  $\tau_{\mathbb{C}}$ -dense ( $\zeta l_{\mathbb{C}}(A) = Y$ ) for every non-empty open subset  $A$  of  $Y$ .
2.  $\mathbb{C}^*$ -hyperconnected if  $Y - \zeta l_{\mathbb{C}}(A) \notin \mathbb{C}$  for every non-empty open subset  $A$  of  $Y$ .
3.  $\mathbb{C}^* \alpha$ -hyperconnected if  $Y - \zeta l_{\mathbb{C}}(A) \notin \mathbb{C}$  for every non-empty  $\mathbb{C}^* \alpha$ -open subset  $A$  of  $Y$ .
4.  $\mathbb{C}^* \alpha o$ -hyperconnected if  $\zeta l_{\mathbb{C}}(A) = Y$  for all  $A \in \mathbb{C}^* \alpha o(Y)$ .

**Proposition 4.2.**

1. Every  $\mathbb{G}^* \alpha$ -hyperconnected is  $*$ -hyperconnected.
2. Every  $*$ -hyperconnected is  $\mathbb{G}^*$ -hyperconnected.
3. Every  $\mathbb{G}^* \alpha$ -hyperconnected is  $\mathbb{G}^*$ -hyperconnected.
4. Every  $\mathbb{G}^* \alpha$ -hyperconnected is  $\mathbb{G}^* \alpha$ -hyperconnected.

**Proof.**


1. Let  $\mathcal{A}$  is a  $\mathbb{G}^* \alpha$ -hyperconnected, then  $cl_{\mathbb{G}}(\mathcal{A}) = \mathbb{Y}$  then  $\mathcal{A}$  is a  $*$ -hyperconnected (since  $\mathcal{A} \in \mathbb{G}^* \alpha(\mathbb{Y})$ . then  $\mathcal{A} \in \tau$ ).

2. Let  $\mathcal{A}$  be a  $*$ -hyperconnected. This means that  $cl_{\mathbb{G}}(\mathcal{A}) = \mathbb{Y}$ . So,  $\mathbb{Y} - cl_{\mathbb{G}}(\mathcal{A}) = \emptyset \notin \mathbb{G}$ . Therefore,  $\mathcal{A}$  is a  $\mathbb{G}^*$ -hyperconnected.

3. Let  $\mathcal{A}$  be an open in  $\mathbb{G}^* \alpha$ -hyperconnected that's mean that  $\mathbb{Y} - cl_{\mathbb{G}}(\mathcal{A}) \notin \mathbb{G}^* \alpha$ -hyperconnected (since every open set in  $\mathbb{G}^* \alpha$ -hyperconnected is an open set in  $\mathbb{G}^*$ -hyperconnected. So,  $\mathbb{Y} - cl_{\mathbb{G}}(\mathcal{A}) \notin \mathbb{G}$ . Therefore,  $\mathcal{A}$  is a  $\mathbb{G}^*$ -hyperconnected.

4. Let  $\mathcal{A}$  be an open in  $\mathbb{G}^* \alpha$ -hyperconnected. This means that  $cl_{\mathbb{G}}(\mathcal{A}) = \mathbb{Y}$ . So,  $\mathbb{Y} - cl_{\mathbb{G}}(\mathcal{A}) \notin \mathbb{G}$ . Therefore,  $\mathcal{A}$  is a  $\mathbb{G}^*$ -hyperconnected.

The following diagram shows the relationship between the types of hyperconnected.



**Diagram 1**  
hyperconnected space via grill space

**4. Conclusion**

In this research we studied a connect space in the grill  $\alpha$  -open topological spaces, we showed some examples and applied some theorems for this new sets. We also found some new properties of these sets.

## References

1. Choquet. G. Sur les notions de filtre et grille, *Comptes Rendus Acad. Sci. Paris*, 224 **1947**, 171-173.
2. Esmaeel, R.B.; Mohammad. R.J On nano soft  $J$ -semi- $g$ -closed sets, *J. Phys. Conf. Ser.* **2020**, 159(1), 012071.
3. Esmaeel, R. B; Nasir. A. I; Bayda Atiya Kalaf. on  $\alpha$ - $\tilde{I}$ -closed soft sets. *Sci. Inter. (Lahore)*. **2018**. 30(5): 703-705.
4. Levine, N. semi-open sets and semi-continuity in topological spaces, *Amer Math. Monthly*, **1963**,70, 36-41.
5. Ahmed Al-Omary ; Takasi Noiri. Decompositions of continuity via grills, *Jordan Journal Mathematics and Statistics (JJMS)* ,**2011**,4(1), 33-46
6. Dhanabal Saravanakumar ; Nagarajan Kalaivani. on grill  $Sp$ - open set in grill topological spaces, *journal of new theory*, **2018**, 23,85-92.
7. Mustafa, M.O.; Esmaeel , R.B. . separation Axioms with Grill-Topological open set, *J. Phys.* **2021**, 1879,2, 022107.
8. Arkhangel'skii, A.V. ; Ponomar"ev, V.I. Fundamentals of general topology-problems and exercises, *Hindustan Pub.Corporation, Delhi*, **1966**.
9. B. Roy ; M. N. Mukherjee. on a typical topology induced by a grill, *Soochow J. Math.*, **2007**, 33 (4) , 771-786.
10. Saad, S. Suliman ; R. B. Esmaeel. On some topology concepts via grill, *Int. J. Nonlinear Anal.* **2022**, Appl. 13 , 1, 3765 – 3772.