



## Effect of Superficially Porous Particles on Chromatographic Analysis Time of Flavonoids

Raid Abdulmehdi Alfaty<sup>1\*</sup>  and Ashraf Saad Rasheed<sup>2</sup> 

<sup>1,2</sup>Department of Chemistry, College of Science, University of Baghdad, Baghdad, Iraq.

\*Corresponding Author.

Received : 1 July 2023

Accepted: 27 August 2023

Published: 20 April 2025

[doi.org/10.30526/38.2.3639](https://doi.org/10.30526/38.2.3639)

### Abstract

Due to superficial particles and remarkable packing technologies, the HALO-HILIC column outperformed regular porous columns in terms of analysis time. This study analyzes and identifies kaempferol and luteolin flavonoids in natural herbs. A fast, novel chromatographic method combining superficially porous silica particles packed in a column served as the separation tool for hydrophilic interaction liquid chromatography coupled with a UV detection system is being introduced in this work. Flavonoid separation and quantification took six minutes using only superficial porous particles in the HILIC column compared to hours using fully porous particles in the RP columns. The findings demonstrated that the HILIC mode might be utilized to determine kaempferol and luteolin levels in ginkgo plant samples. A commercial HALO-HILIC column was used to create the calibration curve, which had the following specifications: linear range ( $0.065\text{--}35\text{ }\mu\text{g mL}^{-1}$  for kaempferol and  $0.065\text{--}9\text{ }\mu\text{g mL}^{-1}$  for luteolin), RSD% not exceeding 0.54%, the limit of quantification ( $0.036\text{ }\mu\text{g mL}^{-1}$  for kaempferol and  $0.024\text{ }\mu\text{g mL}^{-1}$  for luteolin), and limit of determination ( $0.012\text{ }\mu\text{g mL}^{-1}$  for kaempferol and  $0.008\text{ }\mu\text{g mL}^{-1}$  for luteolin).

**Keywords:** HALO-HILIC column, Kaempferol, Luteolin, Medicinal herbs, Superficial particles.

### 1. Introduction

Flavonoids and their derivatives are a significant group of natural compounds found inside cells or outside various plant organs (1). They are primarily sourced from foods like fruits, beverages, herbs, vegetables, and supplements. Natural flavonoids in fruits and vegetables help protect against oxidative stress and serve a nutritional role (2). They are incredibly potent antioxidants and may even be more effective than other now-recognized antioxidants, such as vitamins C and E (3-5). Flavonoids exhibit antioxidant action by preventing free radical synthesis and scavenging ROS, RNS, and other reactive species. Their antioxidant properties are due to the phenolic hydrogen, unique substitution patterns, and chemical composition (6, 7).



Flavonoids comprised of two benzene rings (A and B) joined by a heterocyclic ring (C) through oxygen. Based on the relationship between the B and C rings, the configuration of the B ring, and the hydroxylation and glycosylation sequence of the three rings, flavonoids can be classified into various subclasses (8-10). Anthocyanidins (malvidin, pelargonidin, and peonidin), flavanones (hesperidin, naringenin, and naringin), isoflavones (genistein and daidzein), flavonols (kaempferol, fisetin, and quercetin), and flavones (apigenin, luteolin, and chrysin) are some of the subclasses (11-15). Kaempferol (3,4',5,7-tetrahydroxyflavone) is a flavonoid structurally similar to estrogen, found in various fruits and vegetables like apples, grapes, tomatoes, and green tea (17-19). It has antioxidant, anti-inflammatory, antibacterial, cardiovascular, and neuroprotective properties. Due to its estrogen-like structure, kaempferol may help treat hormone-regulated tumors, including ovarian, breast, cervical, liver cancers, and leukemia (20-22). Luteolin (3',4',5,7-tetrahydroxyflavone), a common flavonoid in fruits, vegetables, and medicinal herbs, is used in Chinese traditional medicine to treat cancer, inflammation, and hypertension (23-26). It acts as an antioxidant or pro-oxidant with anti-inflammatory, anti-allergic, and anti-cancer properties. Recent studies suggest its role in cancer prevention (27-31). The separation and determination of flavonoids in natural sources have increased due to their chemical and biological importance. A common method for isolating and quantifying flavonoids is reversed phase-high-performance liquid chromatography, using various stationary phases and eluents. The mobile phase is often made up of combinations of water and methanol or acetonitrile combinations, while the stationary phases are typically C18 or C8 (32). Even when high-resolution columns are used, co-elution remains a significant issue in studying polyphenolic substances. A further drawback of the reversed-phase mode is the longest single run, which might last between 30 minutes and an hour (33). Separating the flavonoids that have been isolated from nuts and fruit was demonstrated by Harly et al., using a mobile phase consisting of a mixture of methanol, acetonitrile, and trifluoroacetic acid. In under 60 minutes, the authors separated 20 polyphenolic chemicals (34,35). Sakakibara et al., (36) developed a technique for separating and quantifying polyphenolic compounds in food, though it took up to 95 minutes per run. Lee et al., (37) analyzed chokeberry juice, with a single analysis taking 70 minutes. A gradient elution in another study took 65 minutes to determine the polyphenolic composition of Poligano purple carrots (38). Utilizing ultra-pressure liquid chromatography is one method of reducing the duration of a single analysis. This method's foundation is the employment of columns that are densely packed with particles with a diameter of less than 2  $\mu\text{m}$ . In contrast to the usual approach, these findings create high pressure but are distinguished by improved resolution, sensitivity, and retention durations (39-42). Schwarz et al., (43) used UHPLC and traditional HPLC to separate phenolic components in brandy, with UHPLC completing the separation in 6.5 minutes, compared to 60 minutes with HPLC (44). The development of superficially porous particles offers similar efficiencies to UHPLC without high pressure requirements, allowing use with existing HPLC systems, providing financial advantages over UHPLC (45). The column used in this study, introduced in 2006, sparked renewed interest in particle design research. Additionally, hydrophilic interaction liquid chromatography (HILIC) is used as an alternative to RP and NP modes for fast separations (46). In 1990, Alpert first developed the concept of hydrophilic interaction liquid chromatography (47). HILIC typically incorporates polar stationary phases with eluents and many organic solvents (10, 48, 49). Jandera and Janas's study (50) identifies three HILIC

stationary phase subgroups: neutral (e.g., diol), charged (e.g., bare silica), and zwitterionic (e.g., sulfobetaine). The HILIC used in this study is charged bare silica with a fused core and porous shell, offering lower backpressure compared to sub-2- $\mu$ m particles in other columns (51). This study analyzes and identifies kaempferol and luteolin flavonoids in natural herbs.

## 2. Materials and Methods

### 2.1. Apparatus and chemicals

A quaternary pump, a column oven, and a UV-Vis detector were all parts of the HPLC apparatus (Metrohm AG, Germany) for the liquid chromatographic studies. The supplier provided acetonitrile ( $\text{CH}_3\text{CN}$ ) and Formic acid ( $\text{CH}_2\text{O}_2$ ) (HPLC grade) for analysis (Spelco). Sodium acetate is supplied from (Sigma-Aldrich). The water for the mobile phases was purified with a Milli-Q system (from Millipore). Standards for luteolin and kaempferol were provided by Carl Roth GmbH (Karlsruhe, Germany). The HALO-HILIC column was purchased from (Advance Material Technology, USA).

### 2.2. Sample preparation

Ginkgo herb was drained, crushed, and ground before being weighed (1 g) and dissolved in 50 ml of acetonitrile and water (70:30 v/v) to extract flavonoids. The final product was then centrifuged for 30 minutes before being filtered through a 0.45  $\mu\text{m}$  membrane filter after 60 minutes of ultrasonic treatment, 25 minutes of the operation, and 30 minutes of centrifugation.

### 2.3. Standard sample preparation

About 10 mg of each compound was dissolved in 100 mL of  $\text{CH}_3\text{CN}$  to prepare a stock solution (100 mg/L) of kaempferol and luteolin. The solution was next diluted to the proper concentration ranges to create calibration curves and passed through a 0.45  $\mu\text{m}$  filter.


### 2.4. Chromatographic conditions

Employing a HALO-HILIC (100 mm x 2.1 mm ID, 2.7  $\mu\text{m}$ ) to carry out the chromatographic separation of plant flavonoids;  $\text{CH}_3\text{CN}$  with  $\text{CH}_2\text{O}_2$  content served as the comparable mobile phase A (organic portion of the mobile phase), while water with  $\text{CH}_2\text{O}_2$  content served as the comparable mobile phase B (aqueous portion of the mobile phase). The linear gradient elution procedure was established at a 1.0 mL/min flow rate. The column was kept at a constant temperature of 30  $^\circ\text{C}$ , and the injection volume was 10  $\mu\text{L}$ .

## 3. Results and Discussion

### 3.1. Chromatographic separation

In contrast to columns filled with 5  $\mu\text{m}$  porous silica particles, 2.7  $\mu\text{m}$  superficially porous silica particles allow for speedy and efficient HILIC separations. Rapid separations and quantification of highly and moderately polar substances are possible because bare silica columns packed with 2.7  $\mu\text{m}$  particles can be operated at conventional HPLC pressures, and flow rates equal those used in UPLC on RP columns (52). **Figure 1** demonstrates the chromatographic separation of kaempferol and luteolin in ginkgo herbs using a HALO-HILIC commercial column.




**Figure 1.** Chromatographic separation of kaempferol and luteolin in ginkgo herbs using a HALO-HILIC.

It clearly shows the fast run time of the current study (under 6 min) compared to other reported analysis times under the fully poured RP column. The main reason is the particle size and shape of the column used in this study. The effect of superficial particles on the chromatographic behavior best explain by the Van Deemter equation:


$$H = A + \frac{B}{u} + Cu \quad (1)$$

Where H is the plate height (as a measure of the resolving power of a column), u is the velocity; A is the Eddy diffusion (related to channeling through non-ideal packing, B is the longitudinal diffusion (eluting particles in the longitudinal direction resulting dispersion) and C the resistance to mass transfer (resistance to mass transfer coefficient of the analyte between mobile and stationary phase). A, B and C are directly connected to the parameters of the column. Both packing quality and particle size have an impact on eddy diffusion. Compared to fully porous particles, the size distribution of a superficial particle is substantially narrower. By decreasing Eddy diffusion, the distance between the particles in the column is lowered, and efficiency increases. Two significant factors impact A, the first associated with the typical particle packing structure, and the second on homogeneity within the column. A better packing of the spheres was achieved in the case of superficial particles due to the restricted particle distribution (53), as shown in **Figure 2**.



**Figure 2.** The contrast between fully porous particle distribution (traditional silica) and narrow particle distribution (superficial silica) is an example of the impact on peak efficiency and analysis time.

The B term describes axial molecule diffusion during the separation process. Because a core prevents a solute from diffusing through superficially porous particles, a silica column with superficial silica packing diffuses a solute less than one with total silica porosity. As a result, in a Van Deemter equation, the B term decreases in the core-shell superficial silica column, causing faster analysis time (54), as demonstrated in **Figure 3**.




**Figure 3.** An illustration of the effect of the B term is provided by solid core particles preventing diffusion as a solute diffuses through a pore and outside of totally porous particles.

### 3.2. Optimization of chromatographic conditions

#### 3.2.1. Mobile phase influence on the retention factor


The hydrophilicity factor plays a significant part in the behavioral differences between kaempferol and luteolin. Both kaempferol and luteolin exhibit RP activity for commercial columns (HALO-HILIC 2.7  $\mu\text{m}$ ), as shown in **Figure 4**, due to their log  $P_{ow}$  values (2.188 and 2.403), respectively.



**Figure 4.** Effect of acetonitrile percentage on the behavior of kaempferol and luteolin using a HALO-HILIC.

### 3.2.2. Eluent concentration effect on the retention factor

With a rise in eluent concentration, solute behavior has become more hydrophilic, which has resulted in the deactivation of intermolecular ion pairs. Even if  $\text{CH}_3\text{CN}$  exists, this method improves the linearization of the functional column groups. Surprisingly, when the quantity of  $\text{CH}_2\text{O}_2$  is increased from 10-100 mM while retaining  $\text{CH}_3\text{CN}$  at 90% and pH at 5.5, kaempferol and luteolin exhibit enhanced retention factors (**Figure 5**). But at least the hydrophilicity of the analytes may be demonstrated. Additionally, the chromatographic separations are influenced by how well the mobile and pseudo-stationary phases are separated. Creating a pseudo-stationary water layer on the column is believed to be the primary basis for separating kaempferol and luteolin.




**Figure 5.** Effect of eluent concentration on the behavior of kaempferol and luteolin using a HALO-HILIC.

### 3.2.3. Effect of pH on the retention factor

For the separation of kaempferol and luteolin in HILIC mode to be successful, the eluent pH must be altered. The pH increased from 3 to 5.5 using acetic acid/sodium acetate buffer while the eluent concentration was constant at 35 mM and 90%  $\text{CH}_3\text{CN}$ . **Figure 6** demonstrates the rising retention factor of kaempferol and luteolin. The hydroxyl group in the examined flavonoids has been deprotonated in light of the physicochemical information that can be predicted for both flavonoids, which have a  $\text{pK}_a$  value of 7.3 and 7.82, respectively. The analytes' deprotonation is unquestionably discernible when the mobile phase's pH is raised to 5,

causing an increase in the electrostatic interaction between the stationary phase and the analyte and increasing the retention factor.




**Figure 6.** Effect of pH on the behavior of kaempferol and luteolin using a HALO-HILIC.

### 3.3. Linearity and range

In this work, the suggested method's linearity was examined by graphing peak areas against the concentrations of kaempferol and luteolin. The peak area concerning the concentrations of the respective flavonoids was linear in a concentration range of (0.065-35  $\mu\text{g mL}^{-1}$  for kaempferol) and (0.065-9  $\mu\text{g mL}^{-1}$  for luteolin); **Figure 7** shows the Calibration Curve of Kaempferol and Luteolin. **Table 1** shows the linearity of kaempferol and luteolin, regression equation, and calibration statistics. The results show that the peak area and concentration are strongly correlated.

**Table 1.** Linearity, regression, LOD, and LOQ.

| Parameter                           | Flavonoids | HALO-HILIC column |
|-------------------------------------|------------|-------------------|
| Linearity ( $\mu\text{g mL}^{-1}$ ) | Kaempferol | 0.065-35          |
|                                     | Luteolin   | 0.065-9           |
| $R^2$                               | Kaempferol | 0.9997            |
|                                     | Luteolin   | 0.9999            |
| LOD ( $\mu\text{g mL}^{-1}$ )       | Kaempferol | 0.012             |
|                                     | Luteolin   | 0.008             |
| LOQ ( $\mu\text{g mL}^{-1}$ )       | Kaempferol | 0.036             |
|                                     | Luteolin   | 0.024             |



**Figure 7.** Calibration curves of standard solutions of kaempferol and luteolin.

### 3.4. Statistical analysis

Measurements of accuracy, RSD%, and recovery percentage were made on both the same day and different days. This method's effectiveness in **Table 2** is explained by its high recovery values.

**Table 2.** Accuracy and precision.

| Flavonoids | Taken<br>µg/mL | HALO-HILIC |         |           |         |
|------------|----------------|------------|---------|-----------|---------|
|            |                | Inter-Day  |         | Intra-Day |         |
|            |                | Rec. (%)   | RSD (%) | Rec. (%)  | RSD (%) |
| Kaempfero  | 3              | 100.60     | 0.28    | 101.00    | 0.36    |
|            | 4              | 99.00      | 0.41    | 99.50     | 0.45    |
|            | 5              | 100.60     | 0.33    | 100.20    | 0.37    |
| Luteolin   | 3              | 99.20      | 0.52    | 99.00     | 0.39    |
|            | 4              | 100.57     | 0.44    | 100.71    | 0.54    |
|            | 5              | 100.50     | 0.19    | 100.12    | 0.29    |

The results of the t-test and variance ratio F-test methods were used in statistical analyses with a 95% confidence level, as shown in **Table 3**.

**Table 3.** compares the proposed methods 2D with a standard procedure for flavonoid analysis by examining t- and F-statistical tests.

| Flavonoids | Taken<br>µg/mL | Proposed<br>method | Standard method<br>(55) | t-Test<br>(Theor.) | F-Test<br>(Theor.) |
|------------|----------------|--------------------|-------------------------|--------------------|--------------------|
| Kaempferol | 3              | 100.60             | 98.67                   | 0.9085<br>(2.7764) | 1.5790<br>(19.000) |
|            | 4              | 99.00              | 100.13                  |                    |                    |
|            | 5              | 99.00              | 99.55                   |                    |                    |
| Luteolin   | 3              | 99.20              | 100.22                  | 0.4122<br>(2.7764) | 1.9718<br>(19.000) |
|            | 4              | 100.57             | 99.34                   |                    |                    |
|            | 5              | 100.50             | 99.21                   |                    |                    |

These results were compared with those obtained using the standard procedure to assess the effectiveness and competency of the HALO-HILIC approach compared to the standard method

(24). The estimated T and F values were within the theoretical values, indicating little difference in the two approaches' flavonoid quantification accuracy.

### 3.5. Kaempferol and Luteolin content in ginkgo samples

The investigation findings are reported in **Table 4** and show that the proposed approach, which used the HALO- HILIC column, successfully identified and quantified kaempferol and luteolin in black and ginkgo herbs samples.

**Table 4.** Flavonoids contents in ginkgo samples.

| Herbs sample | Flavonoids                      |                               |
|--------------|---------------------------------|-------------------------------|
|              | Kaempferol (mg/g <sup>a</sup> ) | Luteolin (mg/g <sup>a</sup> ) |
| Ginkgo       | 0.325 ± 0.065                   | 0.245 ± 0.021                 |

## 4. Conclusion

The newly developed approach was successfully used to establish the concentrations of kaempferol and luteolin in ginkgo plants. The study also covered the method of collecting and examining the components of these herbs. The analytical process demonstrated great precision, sensitivity, reproducibility, feasibility, and practicability of these technological advancements. This is the outcome of using the HALO-HILIC column. It addressed problems that caused the separation process to be slow when utilizing conventional columns. Based on the method created, the newly established optimization was made.

## Acknowledgment

The authors thank the Department of Chemistry at the College of Science/ University of Baghdad for their assistance.

## Conflict of Interest

There is no conflict of interest in this research

## Funding

There was no financial support for this work.

## Ethical Clearance

The local Scientific Committee at the College of Science/ University of Baghdad approved this study.

## References

1. Grotewold E. The science of flavonoids. Springer, 2006. <https://link.springer.com/book/10.1007/978-0-387-28822-2>.
2. Holiman PC, Hertog MG, Katan MB. Analysis and health effects of flavonoids. Food Chem. 1996; 57(1):43-46. [https://doi.org/10.1016/0308-8146\(96\)00065-9](https://doi.org/10.1016/0308-8146(96)00065-9).
3. Chen ZY, Chan PT, Ho KY, Fung KP, Wang J. Antioxidant activity of natural flavonoids is governed by number and location of their aromatic hydroxyl groups. Chem Phys Lipids. 1996; 79(2): 157-163. [https://doi.org/10.1016/0009-3084\(96\)02523-6](https://doi.org/10.1016/0009-3084(96)02523-6).
4. Kopustinskiene DM, Jakstas V, Savickas A, Bernatoniene J. Flavonoids as anticancer agents. Nutrients. 2020; 12(2):457. <https://doi.org/10.3390/nu12020457>.

5. Russo M, Moccia S, Spagnuolo C, Tedesco I, Russo G L, Roles of flavonoids against coronavirus infection. *Chem Biol Interact.* 2020; 328, 109211. <https://doi.org/10.1016/j.cbi.2020.109211>.
6. Chen GL, Fan MX, Wu JL, Li N, Guo MQ. Antioxidant and anti-inflammatory properties of flavonoids from lotus plumule. *Food Chem.* 2019; 277:706-712. <https://doi.org/10.1016/j.foodchem.2018.11.040>.
7. Ninfali P, Antonelli A, Magnani M, Scarpa ES. Antiviral properties of flavonoids and delivery strategies. *Nutrients.* 2020; 12(9):2534. <https://doi.org/10.3390/nu12092534>.
8. Wang TY, Li Q, Bi KS. Bioactive flavonoids in medicinal plants: Structure, activity and biological fate. *AJPS.* 2018; 13(1):12-23. <https://doi.org/10.1016/j.ajps.2017.08.004>.
9. Teng H, Deng H, Zhang C, Cao H, Huang Q, Chen L. The role of flavonoids in mitigating food originated heterocyclic aromatic amines that concerns human wellness. *Food Sci Hum. Wellness.* 2023; 12(4):975-985. <https://doi.org/10.1016/j.fshw.2022.10.019>.
10. Waheb AA, Rasheed AS, Hassan MJM. Quantification of ketoprofen in film-coated tablets using hydrophilic interaction liquid chromatography-Ultraviolet spectrometry. *EJCHEM.* 2022; 65(9):1-4. <https://doi.org/10.21608/ejchem.2022.113077.5138>.
11. Kumar S, Pandey AK. Chemistry and biological activities of flavonoids: an overview. *Sci World J.* 2013; 62750:1-16. <https://doi.org/10.1155/2013/162750>.
12. Kr RR, Rasheed AS, Hassan MJM. Zwitterion chromatography-hydrophilic interaction chromatography for separation and quantitative of rutin and quercetin from herbs and bee products. *J Chem Soc Pak.* 2021; 43(4):484-492.
13. AL-Ayash AS, Khammas ZA, Jasim F. Determination of desferrioxamine in the drug desferal™ as dfom-Au (iii) complex by using indirect electrothermal atomic absorption spectrometry and other techniques. *Baghdad Sci J.* 2008; 5(3):409-415. <https://doi.org/10.21123/bsj.2008.11874>.
14. Teng H, Zheng Y, Cao H, Huang Q, Xiao J, Chen L. Enhancement of bioavailability and bioactivity of diet-derived flavonoids by application of nanotechnology: A review. *Crit Rev Food Sci Nutr.* 2023; 63(3):378-393. <https://doi.org/10.1080/10408398.2021.1947772>.
15. Shi J, Yang G, You Q, Sun S, Chen R, Lin Z, Simal-Gandara J, Lv H. Updates on the chemistry, processing characteristics, and utilization of tea flavonoids in last two decades (2001-2021). *Crit Rev Food Sci Nutr.* 2023; 63(20):4757-4784. <https://doi.org/10.1080/10408398.2021.2007353>.
16. Pandey KB, Rizvi SI. Plant polyphenols as dietary antioxidants in human health and disease. *Oxid Med Cell Longev.* 2009; 2(5):270-278. <https://doi.org/10.4161/oxim.2.5.9498>.
17. Yang L, Gao Y, Bajpai VK, El-Kammar HA, Simal-Gandara J, Cao H, Cheng KW, Wang M, Arroo RR, Zou L, Farag MA, Zhao Y, Xiao J. Advance toward isolation, extraction, metabolism and health benefits of kaempferol, a major dietary flavonoid with future perspectives. *Crit Rev Food Sci Nutr.* 2023; 63(16):2773-2789. <https://doi.org/10.1080/10408398.2021.1980762>.
18. Bian Y, Lei J, Zhong J, Wang B, Wan Y, Li J, Liao C, He Y, Liu Z, Ito K, Zhang B. Kaempferol reduces obesity, prevents intestinal inflammation, and modulates gut microbiota in high-fat diet mice. *J Nutr Biochem.* 2022; 99:108840. <https://doi.org/10.1016/j.jnutbio.2021.108840>.
19. Alam W, Khan H, Shah MA, Cauli O, Saso L. Kaempferol as a dietary anti-inflammatory agent: current therapeutic standing. *Molecules.* 2020; 25(18):4073. <https://doi.org/10.3390/molecules25184073>.
20. Imran M, Salehi B, Sharifi-Rad J, Gondal TA, Saeed F, Imran A, Shahbaz M, Fokou PVT, Arshad MU, Khan H, Guerreiro SG, Martins N, Estevinho LM. Kaempferol: A key emphasis to its anticancer potential. *Molecules.* 2019; 24(12):2277. <https://doi.org/10.3390/molecules24122277>.

21. Alshehri AS. Kaempferol attenuates diabetic nephropathy in streptozotocin-induced diabetic rats by a hypoglycaemic effect and concomitant activation of the Nrf-2/Ho-1/antioxidants axis. *Arch Physiol Biochem*. 2023; 129(4):984-997. <https://doi.org/10.1080/13813455.2021.1890129>.
22. Santos JSD, Cirino JPG, Carvalho PDO, Ortega MM. The pharmacological action of kaempferol in central nervous system diseases: A review. *Front Pharmacol*. 2021; 11:565700. <https://doi.org/10.3389/fphar.2020.565700>.
23. Azeem M, Hanif M, Mahmood K, Ameer N, Chughtai FRS, Abid U. An insight into anticancer, antioxidant, antimicrobial, antidiabetic and anti-inflammatory effects of quercetin: A review. *Polym Bull (Berl)*. 2023; 80(1):241-262. <https://doi.org/10.1007/s00289-022-04091-8>.
24. Caporali S, De Stefano A, Calabrese C, Giovannelli A, Pieri M, Savini I, Tesaro M, Bernardini S, Minieri M, Terrinoni A. Anti-inflammatory and active biological properties of the plant-derived bioactive compounds luteolin and luteolin 7-glucoside. *Nutrients*. 2022; 14(6):1155. <https://doi.org/10.3390/nu14061155>.
25. Franza L, Carusi V, Nucera E, Pandolfi F. Luteolin, inflammation and cancer: Special emphasis on gut microbiota. *Biofactors*. 2021; 47(2):181-189S. <https://doi.org/10.1002/biof.1710>.
26. AL-Ayash AS, Muhamad YH, Ghafouri SA. Spectrophotometric determination of epinephrine in pharmaceutical preparations using praseodymium as mediating metals. *Baghdad Sci J*. 2011; 8(1): 110-117. <https://doi.org/10.21123/bsj.2011.8.1.110-117>.
27. Ge X, He X, Liu J, Zeng F, Chen L, Xu W, Shao R, Huang Y, Farag MA, Capanoglu E, El-Seedi HR, Zhao C, Liu B. Amelioration of type 2 diabetes by the novel 6, 8-guanidyl luteolin quinone-chromium coordination via biochemical mechanisms and gut microbiota interaction. *J Adv Res*. 2023; 46:173-188. <https://doi.org/10.1016/j.jare.2022.06.003>.
28. Nguyen TLA, Bhattacharya D. Antimicrobial activity of quercetin: an approach to its mechanistic principle. *Molecules*. 2022; 27(8):2494. <https://doi.org/10.3390/molecules27082494>.
29. Di Stadio A, D'Ascanio L, Vaira LA, Cantone E, De Luca P, Cingolani C, Motta G, De Riu G, Vitelli F, Spriano G, De Vincentiis M, Camaioni A, La Mantia I, Ferrel F, Brenner MJ. Ultramicrosized palmitoylethanolamide and luteolin supplement combined with olfactory training to treat post-COVID-19 olfactory impairment: a multi-center double-blinded randomized placebo-controlled clinical trial. *Curr Neuropharmacol*. 2022; 20(10):2001-2012. <https://doi.org/10.2174/1570159x20666220420113513>.
30. Theoharides TC, Cholevas C, Polyzoidis K, Politis A. Long-COVID syndrome-associated brain fog and chemofog: Luteolin to the rescue. *Biofactors*. 2021; 47(2):232-241. <https://doi.org/10.1002/biof.1726>.
31. ChemAxon EU UGM. Predictive data. 2022.
32. Merken HM, Beecher GR. Measurement of food flavonoids by high-performance liquid chromatography: a review. *J Agric Food Chem*. 2000; 48 (3):577-599. <https://doi.org/10.1021/jf990872o>.
33. Singh S, Uddin M, Khan MMA, Chishti AS, Bhat UH. Therapeutic properties of capsaicin: a medicinally important bio-active constituent of chilli pepper. *AJPCR*. 2022; 15(7):47-58. <http://dx.doi.org/10.22159/ajpcr.2022.v15i7.44405>.
34. Harnly JM, Doherty RF, Beecher GR, Holden JM, Haytowitz DB, Bhagwat S, Gebhardt S. Flavonoid content of US fruits, vegetables, and nuts. *J Agric Food Chem*. 2006; 54(26):9966-9977. <https://doi.org/10.1021/jf061478a>.
35. AL-Ayash AS, Jasim F, Zair T. Spectrophotometric micro determination of drug promethazine hydrochloride in some pharmaceutical by chelating with Rhodium. *Baghdad Sci J*. 2008; 5(4):639-645. <https://doi.org/10.21123/bsj.2008.5.4.638-645>

36. Sakakibara H, Honda Y, Nakagawa S, Ashida H, Kanazawa K. Simultaneous determination of all polyphenols in vegetables, fruits, and teas. *J Agric Food Chem.* 2003; 51(3):571-581. <https://doi.org/10.1021/jf020926l>.
37. Lee JE, Kim GS, Park S, Kim YH, Kim MB, Lee WS, Jeong SW, Lee SJ, Jin JS, Shin SC. Determination of chokeberry (*Aronia melanocarpa*) polyphenol components using liquid chromatography–tandem mass spectrometry: Overall contribution to antioxidant activity. *Food Chem.* 2014; 146:1-5. <https://doi.org/10.1016/j.foodchem.2013.09.029>.
38. Pace B, Capotorto I, Cefola M, Minasi P, Montemurro N, Carbone V. Evaluation of quality, phenolic and carotenoid composition of fresh-cut purple Polignano carrots stored in modified atmosphere. *J Food Compos Anal.* 2020; 86:103363. <https://doi.org/10.1016/j.jfca.2019.103363>.
39. Jing J, Parekh HS, Wei M, Ren WC, Chen SB. Advances in analytical technologies to evaluate the quality of traditional Chinese medicines. *Trac-Trend Anal Chem.* 2013; 44:39-45. <https://doi.org/10.1016/j.trac.2012.11.006>.
40. Fahad AMM, Rasheed AS, Ali HH. Hydrophilic interaction chromatography with sulfobetaine zwitterionic polymer-bonded stationary phases for the simultaneous quantification of atorvastatin and rosuvastatin pharmaceuticals in bulk and dosage forms. *JCSP.* 2022; 44(4):330-337. <https://doi.org/10.52568/001070/JCSP/44.04.2022>.
41. Fahad AMM, Rasheed AS, Ali HH. Separation and determination of simvastatin on ZIC-HILIC stationary phases by hydrophilic interaction chromatography in pharmaceutical material products. *Mater Today: Proc.* 2022; 49 (7):2817-2821. <https://doi.org/10.1016/j.matpr.2021.09.535>.
42. Qassim AW, Rasheed AS, Abdulrahman SK. Hydrophilic interaction chromatographic analysis of genistein in herbs and propolis. *Sys Rev Pharm.* 2020; 11(12):756-763.
43. Schwarz M, Rodríguez MC, Guillén DA, Barroso CG. Development and validation of UPLC for the determination of phenolic compounds and furanic derivatives in Brandy de Jerez. *JSS.* 2009; 32(11): 1782-1790. <https://doi.org/10.1002/jssc.200800706>.
44. Salisbury JJ. Fused-core particles: a practical alternative to sub-2 micron particles. *JCS.* 2008; 46(10):883-886. <https://doi.org/10.1093/chromsci/46.10.883>.
45. Ali I, AL-Othman ZA, Al-Za'abi M. Superficially porous particles columns for super fast HPLC separations. *BMC.* 2012; 26(8):1001-1008. <https://doi.org/10.1002/bmc.2690>.
46. Fadhil AK, Rasheed AS, Hassan MJM. A review of recent advances in the estimation of pharmaceutical products using hydrophilic interaction chromatography (HILIC) technology. *Egypt. J. Chem.* 2023; 66(1):399-417. <https://doi.org/10.21608/ejchem.2022.121396.5443>.
47. Alpert AJ. Hydrophilic-interaction chromatography for the separation of peptides, nucleic acids and other polar compounds. *J Chromatogr A.* 1990; 499:177-196. [https://doi.org/10.1016/S0021-9673\(00\)96972-3](https://doi.org/10.1016/S0021-9673(00)96972-3).
48. Waheb AA, Rasheed AS, Hassan MJM. Strategies for the separation and quantification of non-steroidal anti-inflammatory drugs using ZIC-HILIC-HPLC with UV detection. *Curr Pharm Anal.* 2022; 18(10):949-958. <https://doi.org/10.2174/1573412918666220915090831>.
49. Fadhil AK, Rasheed AS, Hassan MJM. Evaluation and application of ZIC-HILIC columns selectivity for four angiotensin II receptor blockers in pharmaceutical formulations. *Curr Pharm Anal.* 2022; 18 (9):901-908. <https://dx.doi.org/10.2139/ssrn.4008001>.
50. Jandera P, Janás P. Recent advances in stationary phases and understanding of retention in hydrophilic interaction chromatography. A review. *Anal Chim Acta.* 2017; 967:12-32. <https://doi.org/10.1016/j.aca.2017.01.060>.
51. Chauve B, Guilleme D, Cléon P, Veuthey JL. Evaluation of various HILIC materials for the fast separation of polar compounds. *J Sep Sci.* 2010; 33(6-7):752-764. <https://doi.org/10.1002/jssc.200900749>.

52. Hsieh Y, Galviz G, Long BJ. Ultra-performance hydrophilic interaction liquid chromatography/tandem mass spectrometry for the determination of everolimus in mouse plasma. *Rapid Commun Mass Spectrom.* 2009; 23(10):1461-1466. <https://doi.org/10.1002/rcm.4022>.
53. Gritti F, Guiochon G. Diffusion models in chromatographic columns packed with fully and superficially porous particles. *Chem Eng Sci.* 2011; 66(17):3773-3781. <https://doi.org/10.1016/j.ces.2011.04.039>.
54. Gumustas M, Zalewski P, Ozkan SA, Uslu B. The history of the core-shell particles and applications in active pharmaceutical ingredients via liquid chromatography. *Chromatographia.* 2019; 82:17-48. <https://doi.org/10.1007/s10337-018-3670-6>.
55. Collison MW. Determination of total soy isoflavones in dietary supplements, supplement ingredients, and soy foods by high-performance liquid chromatography with ultraviolet detection: collaborative study. *Journal of AOAC International.* 2008; 91(3):489-500.