



A Numerical Study of Initial and Boundary Conditions Problem for Wave Equation Using the Operational Matrices

Myasar Obaid Enadi^{1*} and Majeed A. AL-Jawary²

^{1,2}Department of Mathematics, College of Education for Pure Sciences (Ibn AL-Haitham), University of Baghdad, Baghdad, Iraq.

*Corresponding Author.

Received: 6 December 2024

Accepted: 12 May 2025

Published: 20 October 2025

doi.org/10.30526/38.4.4097

Abstract

In this paper, orthogonal polynomials and their operational matrices will be utilized to address the initial and boundary value problems of the one-dimensional wave problem, where the domain of the space variable is bounded, which covers a variety of scientific and engineering operations. Six types of orthogonal polynomials, Include instead of such as the Genocchi, Bernoulli, Legendre, Boubaker, Chebyshev and Standard polynomials. The linear problem with its initial and boundary conditions are transformed to a linear algebraic equations, which can then be solved by utilizing Mathematica[®]12 to get an approximate solution for this problem. Some test problems related to the one-dimensional wave equation with different conditions are discussed and solved to show how reliable and efficient the proposed methods. The error norm L_{∞} and the mean square error L_{rms} , were computed; these are presented through analytical tables and graphics showing the rapid convergence for these methods.

Keywords: Wave problem; Operational matrices, Orthogonal Polynomials, Approximate solutions.

1. Introduction

The mathematical foundation for many problems in mathematics, physics, engineering, and chemistry can be explained mathematically using partial differential equations (PDEs). In physics, for example, partial differential equations clear describe wave propagation and heat transfer. Furthermore, partial differential equations were utilized to describe the majority of physical processes in domains such as electricity, quantum mechanics, wave propagation in shallow water, fluid dynamics, plasma physics, and others (1).

There is a need to find effective and reliable approximate or analytical techniques that can handle PDEs because of these massive applications, there is a need to develop effective and trustworthy approximate or analytical procedures for dealing with PDEs. Many mathematicians and engineers have solved a wide range of functional equations using the VIM by He (2–4).

Many tapes of scientific, physical, and engineering problems can be classified as initial boundary value problems (IBVP). With a few exceptions, we are unable to find accurate analytical answers to the majority of these difficulties. There have been several attempts to



develop approximate and analytical methods for solving the non-linear diffusion and wave problems, see (5–8).

Wave equations can only be solved analytically in highly particular situations, hence many realistic cases cannot be solved using commonly used analytical methods. In addition to discretization methods such as finite difference, finite volume, and finite element approaches, various more ways have been proposed to solve the wave problem. For example, the analytical approach to solving the wave equation (9), the numerical solution of the wave issue (10,11), and the optimal homotopy strategy for solving the nonlinear wave equation (12). Furthermore, there are numerous techniques that offer an approximate resolve for the differing kinds of the differential equations, see, (13–15).

Corrington suggested in 1973 that a system of algebraic linear equations could be created from linear differential and integral equations by using a least squares approximation and repeatedly integrating Walsh functions (16). In addition, Sparis and Mouroutsos in 1986 using the orthogonal polynomial series operational matrices to solve differential equations (17), many researchers have used orthogonal polynomials to find approximate solutions for many applications. See; (18–28). The authors were highly interested in since they were practical methods for resolving an extensive number of approximation theory and numerical analysis problems. On the other hand, the orthogonal polynomials and operational matrices stand out above other types due to their effective reduction of the required solution, which is achieved by applying the operational matrices technique to transform the linear differential equations into linear algebraic systems of equations, where any computer program can be used to solve them.

Farther more, Turkyilmazoglu in 2013 proposed an approximate analytical technique for resolving differential equations based on standard polynomials, which is used on various types of problems (29–33).

In 2023, Othman, et al. introduced several orthogonal polynomials, for instance, Hermite, Laguerre, Chebyshev, and others polynomials with inner product, to develop the computational method (CM) (34). Myasar, et al. (35,36), further introduced the Genocchi, Bernoulli, and Boubaker polynomials, which contribute to the effective computational method (ECM).

The outline for this paper is as follows: Section two shows the second-order linear wave equation's mathematical formulation. Section three: Preliminary of the orthogonal polynomials. Section four: Main results and applications of orthogonal polynomials to resolve some examples for the wave problem. Section five: gives the conclusions.

2. The Mathematical Model of the Wave Equation

Let us consider the one-dimensional wave problem (7):

$$\frac{\partial^2 u}{\partial t^2} - c^2 \frac{\partial^2 u}{\partial x^2} = h(x, t), \quad t \geq 0, x \in \Omega. \quad (1)$$

with initial conditions:

$$u(x, 0) = f(x), \quad \frac{\partial u}{\partial t}(x, 0) = g(x), \quad x \in \partial_D \Omega, \quad (2)$$

and boundary conditions:

$$u(x, t) = \hat{u}(x, t), \quad t \geq 0, x \in \partial_D \Omega, \quad (3)$$

$$Fu(x, t) = \hat{f}(x, t), \quad t \geq 0, x \in \partial_N \Omega. \quad (4)$$

where $h(x, t)$ is a particular function, Ω is the bounded domain, c is a constant, $\hat{f}(x, t)$ and $\hat{u}(x, t)$ are known functions, $[Fu](x, t) = \frac{\partial u}{\partial n}(x, t)$, and $n(x, t)$ the external normal vector to the boundary $\partial \Omega$, when $f = 0$ **Equation 1** will be homogeneous wave equation, and if $f \neq 0$ **Equation 1** will be non-homogeneous wave problem.

3. Preliminary of the orthogonal polynomials and their operational matrices

The orthogonal polynomials and their operational matrices play an important part applied and pure mathematics as well as numerical calculation. Six types of these orthogonal polynomials will be used: standard polynomials (x^i), Legendre, Chebyshev, Boubaker, Genocchi, and Bernoulli polynomials to resolve the wave problem with initial and boundary conditions.

3.1. The operational matrices for standard polynomials

Assume that the wave problem (1), with the initial and boundary conditions (2-4), has a unique solution, this can be implemented using a suitable linear transformation. Taking the basic functions

$$\xi = \{\xi_0(x), \xi_1(x), \dots, \xi_n(x), \dots\}, \text{ and } \eta = \{\eta_0(t), \eta_1(t), \dots, \eta_n(t), \dots\},$$

Let the solution of **Equation 1**, may be presented by using the double series extension in terms of a base functions

$$u(x, t) = \sum_{h=0}^{\infty} \sum_{k=0}^{\infty} b_{hk} \xi_h(x) \eta_k(t), \quad (5)$$

The **Equation 5** may be written using the dot product, as follows:

$$u(x, t) = \sum_{h=0}^n \sum_{k=0}^n b_{hk} \xi_h(x) \eta_k(t) = \psi(x) \cdot B \cdot \phi(t), \quad (6)$$

where b_{hk} the coefficients will be evaluated later, by use the definition:

$$\psi(x) = [1 \ x \ \dots \ x^n], \ \phi(t) = [1 \ t \ \dots \ t^n]^T, \text{ and } B = \begin{bmatrix} b_{00} & b_{01} & \dots & b_{0n} \\ b_{10} & b_{11} & \dots & b_{1n} \\ \vdots & \vdots & \ddots & \vdots \\ b_{n0} & b_{n1} & \dots & b_{nn} \end{bmatrix}^T,$$

Furthermore, the n^{th} -order partial derivatives for ψ and ϕ with respect x and t are obtained by:

$$\frac{\partial^n \psi(x)}{\partial x^n} = \psi(x) \cdot A_S, \quad \frac{\partial^n \phi(t)}{\partial t^n} = (A_S)^T \cdot \phi(t),$$

where A_S is the derivative matrix that is dimensional $(n+1) \times (n+1)$ is given by:

$$A_S = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & \ddots & \vdots \\ \vdots & \vdots & \vdots & \dots & n \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix},$$

Thus, the derivatives of $u(x, t)$ will be obtained using **Equation 6** have the following forms:

$$\frac{\partial^n u(x, t)}{\partial x^n} = \psi(x) \cdot A_S \cdot B \cdot \phi(t), \quad \frac{\partial^n u(x, t)}{\partial t^n} = \psi(x) \cdot B \cdot (A_S)^T \cdot \phi(t), \quad n \geq 1. \quad (7)$$

3.2. The operational matrices for Boubaker polynomials

The n^{th} -degree of Boubaker polynomials $P_n(x)$ are defined as follows (28,37):

$$P_n(x) = \sum_{k=0}^{(n/2)} (-1)^k \binom{n-k}{k} \frac{n-4k}{n-k} x^{n-2k},$$

also Boubaker polynomials can be computed by used the iterative procedure as follows:

$$P_{m+1}(x) = x P_m(x) - P_{m-1}(x), \quad m \geq 2.$$

Furthermore, suppose that the unknown function $u(x, t)$, may be approximated by applying the double series based on the Boubaker polynomial:

$$u(x, t) = \sum_{h=0}^n \sum_{k=0}^n b_{hk} P_h(x) P_k(t) = \psi(x) \cdot B \cdot \phi(t), \quad (8)$$

where $\psi(x) = [P_0(x), \dots, P_n(x)]$, $\phi(t) = [P_0(t), \dots, P_n(t)]^T$, and

$$B = \begin{bmatrix} b_{00} & b_{01} & \dots & b_{0n} \\ b_{10} & b_{11} & \dots & b_{1n} \\ \vdots & \vdots & \ddots & \vdots \\ b_{n0} & b_{n1} & \dots & b_{nn} \end{bmatrix}^T,$$

where b_{hk} are the Boubaker polynomials coefficients will be evaluated latter.

The derivatives of $u(x, t)$ may be transformed into matrices by utilizing the following formulation:

$$\frac{\partial^n u(x, t)}{\partial x^n} = \psi(x) \cdot A_{Bk} \cdot B \cdot \phi(t), \quad \frac{\partial^n u(x, t)}{\partial t^n} = \psi(x) \cdot B \cdot (A_{Bk})^T \cdot \phi(t), \quad n \geq 1. \quad (9)$$

The derivatives matrix of Boubaker polynomials denoted by A_{Bk} , and if N is even or N is odd, Defined A_{Bk} as follows, respectively:

$$A_{Bk \text{ even}} = \begin{bmatrix} a_{0,0} & 0 & 0 & \cdots & 0 & 0 & 0 \\ 0 & a_{1,0} & 0 & \cdots & 0 & 0 & 0 \\ a_{2,1} & 0 & a_{2,0} & \cdots & 0 & 0 & 0 \\ 0 & a_{3,1} & 0 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ a_{N-1, \frac{N-1}{2}} & 0 & a_{N-1, \frac{N-3}{2}} & \cdots & 0 & a_{N-1,0} & 0 \\ 0 & a_{N, \frac{N-1}{2}} & 0 & \cdots & a_{N,1} & 0 & a_{N,0} \end{bmatrix},$$

$$A_{Bk \text{ odd}} = \begin{bmatrix} a_{0,0} & 0 & 0 & \cdots & 0 & 0 & 0 \\ 0 & a_{1,0} & 0 & \cdots & 0 & 0 & 0 \\ a_{2,1} & 0 & a_{2,0} & \cdots & 0 & 0 & 0 \\ 0 & a_{3,1} & 0 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & a_{N-1, \frac{N-2}{2}} & 0 & \cdots & 0 & a_{N-1,0} & 0 \\ a_{N, \frac{N}{2}} & 0 & a_{N, \frac{N-2}{2}} & \cdots & a_{N,1} & 0 & a_{N,0} \end{bmatrix}.$$

The following relations can be used to calculate the components $\{a_{n,k}\}_{n=0, k=0}^{N, \lfloor \frac{n}{2} \rfloor}$:

$$a_{n,0} = 1, \quad a_{n,1} = -(n-4), \quad a_{n,k} = (-1)^k \frac{n-4k}{k!} \prod_{j=k+1}^{2k-1} (n-j), \quad k = 2, 3, \dots, \left\lfloor \frac{n}{2} \right\rfloor.$$

3.3. The operational matrices for Bernoulli polynomials

Bernoulli polynomials $B_m(x)$ of m^{th} – degree is given by (27,38,39):

$$B_m(x) = \sum_{j=0}^m \binom{m}{j} B_j x^{m-j},$$

where B_j is the Bernoulli number, such that $B_j = -\sum_{j=0}^{n+1} \frac{(-1)^j}{j} \binom{n+1}{j} \sum_{h=1}^j h^n$, for $n \geq 0$, $n \neq 1$,

The first Bernoulli numbers define as follows:

$$B_0 = 1, B_1 = -\frac{1}{2}, B_2 = \frac{1}{6}.$$

Additionally, the unknown function $u(x, t)$ can be approximated by using the following dot product based on the Bernoulli polynomials:

$$u(x, t) = \sum_{h=0}^n \sum_{k=0}^n b_{hk} P_h(x) P_k(t) = \psi(x) \cdot B \cdot \phi(t), \quad (10)$$

where $\psi(x) = [B_0(x), \dots, B_n(x)]$, $\phi(t) = [B_0(t), \dots, B_n(t)]^T$, and

$$B = \begin{bmatrix} b_{00} & b_{01} & \cdots & b_{0n} \\ b_{10} & b_{11} & \cdots & b_{1n} \\ \vdots & \vdots & \ddots & \vdots \\ b_{n0} & b_{n1} & \cdots & b_{nn} \end{bmatrix}^T,$$

where b_{hk} is the unknown coefficients of Bernoulli polynomials determined latter.

Moreover, the derivatives of $u(x, t)$ can be written in matrices form by the following formula:

$$\frac{\partial^n u(x, t)}{\partial x^n} = \psi(x) \cdot A_S \cdot B \cdot \phi(t), \quad \frac{\partial^n u(x, t)}{\partial t^n} = \psi(x) \cdot B \cdot (A_S)^T \cdot \phi(t), \quad n \geq 1. \quad (11)$$

where A_{Br} is the derivative matrix of Bernoulli polynomials, is given as below:

$$A_{Br} = \begin{bmatrix} 0 & 0 & 0 & \cdots & 0 \\ 1 & 0 & 0 & \cdots & 0 \\ 0 & 2 & 0 & \cdots & 0 \\ \vdots & 0 & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & k & 0 \end{bmatrix},$$

3.4. The operational matrices for Legendre polynomials

Legendre polynomials $L_m(x)$ of m^{th} -degree is given as follows (24,40):

$$L_0(x) = 1, \quad L_1(x) = x, \quad \dots, \quad L_{n+1}(x) = \frac{x(2n+1)L_n(x) - nL_{n-1}(x)}{n+1}, \quad n = 2, 3, \dots$$

Furthermore, the Legendre polynomials will be get by using the formula as follows:

$$L_k(x) = \sum_{i=0}^n (-1)^{n+i} \frac{(n+i)!}{2^i (n-i)! (i!)^2} (x+1)^i.$$

By using, the linear combination as follow can be approximated the unknown function $u(x, t)$.

$$u(x, t) = \sum_{h=0}^n \sum_{k=0}^n b_{hk} L_h(x) L_k(t) = \psi(x). B. \phi(t), \quad (12)$$

where $\psi(x) = [L_0(x), \dots, L_n(x)]$, $\phi(t) = [L_0(t), \dots, L_n(t)]^T$, and

$$B = \begin{bmatrix} b_{00} & b_{01} & \cdots & b_{0n} \\ b_{10} & b_{11} & \cdots & b_{1n} \\ \vdots & \vdots & \ddots & \vdots \\ b_{n0} & b_{n1} & \cdots & b_{nn} \end{bmatrix}^T,$$

Moreover, b_{hk} is the coefficients of Legendre polynomials evaluated latter.

The derivative of $u(x, t)$ with a respect to x and t may be written in matrices as follows:

$$\frac{\partial^n u(x, t)}{\partial x^n} = \psi(x). A_L. B. \phi(t), \quad \frac{\partial^n u(x, t)}{\partial t^n} = \psi(x). B. (A_L)^T. \phi(t), \quad n \geq 1. \quad (13)$$

The derivatives matrix of the Legendre polynomials denoted by A_L , and we obtained by (34):

$$A_L = \begin{cases} (2j+1), & j = i - n, \\ 0 & \text{otherwise.} \end{cases}$$

If k is even then $n = 1, 3, \dots, k-1$, and if k is odd then $n = 1, 3, \dots, k$.

3.5. The operational matrices for Genocchi polynomials

The Genocchi polynomials $G_n(x)$ of n^{th} -degree is given by (20):

$$G(x) = \sum_{j=0}^n \binom{n}{j} G_n x^{n-j},$$

By using, the linear combination as follow can be approximated the function $u(x, t)$.

$$u(x, t) = \sum_{h=0}^n \sum_{k=0}^n b_{hk} G_h(x) G_k(t) = \psi(x). B. \phi(t), \quad (14)$$

where $\phi(t) = [G_0(t), \dots, G_n(t)]^T$, $\psi(x) = (G_0(x), \dots, G_n(x))$ and

$$B = \begin{bmatrix} b_{00} & b_{01} & \cdots & b_{0n} \\ b_{10} & b_{11} & \cdots & b_{1n} \\ \vdots & \vdots & \ddots & \vdots \\ b_{n0} & b_{n1} & \cdots & b_{nn} \end{bmatrix}^T$$

Moreover, b_{hk} is the unknown coefficient of Genocchi polynomials evaluated latter.

The derivatives of $u(x, t)$ can be written in matrices form as follows:

$$\frac{\partial^n u(x, t)}{\partial x^n} = \psi(x). G. B. \phi(t), \quad \frac{\partial^n u(x, t)}{\partial t^n} = \psi(x). B. (G)^T. \phi(t), \quad n \geq 1. \quad (15)$$

The derivative matrix of Genocchi polynomials is denoted by G and given by:

$$G = \begin{bmatrix} 0 & 0 & 0 & \cdots & 0 & 0 \\ 2 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 3 & 0 & \cdots & 0 & 0 \\ 0 & 0 & \ddots & 0 & 0 & 0 \\ \vdots & \vdots & \cdots & n-1 & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & n & 0 \end{bmatrix},$$

3.6. The operational matrices for Chebyshev polynomials

Chebyshev polynomials $H_m(x)$ of m^{th} – degree are defined as follows (25,41):

$$H_m(x) = \sum_{i=0}^m (-1)^{m-i} 2^i \frac{(m+i-1)!}{(m-i)!(2i)!} (x+1)^i,$$

In addition, the function $u(x, t)$ may be approximated by using the linear combination as following:

$$u(x, t) = \sum_{h=0}^n \sum_{k=0}^n b_{hk} H_h(x) H_k(t) = \psi(x) \cdot B \cdot \phi(t), \quad (16)$$

where $\phi(t) = [H_0(t), \dots, H_n(t)]^T$, $\psi(x) = [H_0(x), \dots, H_n(x)]$, and

$$B = \begin{bmatrix} b_{00} & b_{01} & \cdots & b_{0n} \\ b_{10} & b_{11} & \cdots & b_{1n} \\ \vdots & \vdots & \ddots & \vdots \\ b_{n0} & b_{n1} & \cdots & b_{nn} \end{bmatrix}^T,$$

Furthermore, b_{hk} represents the unknown coefficients of Chebyshev polynomials calculated latter.

The derivatives of $u(x, t)$ may be rewritten in matrices using the following formula:

$$\frac{\partial^n u(x, t)}{\partial x^n} = \psi(x) \cdot A_{Ch} \cdot B \cdot \phi(t), \quad \frac{\partial^n u(x, t)}{\partial t^n} = \psi(x) \cdot B \cdot (A_{Ch})^T \cdot \phi(t), \quad n \geq 1. \quad (17)$$

The derivative matrix of the Chebyshev polynomials is denoted by A_G , and obtained by:

$$A_{Ch} = \begin{cases} \frac{2!}{\xi_i} & i = k - m, \\ 0 & \text{otherwise,} \end{cases}$$

If j is odd then $m = 1, 3, \dots, j$ and if j is even then $m = 1, 3, \dots, j - 1$, and $\xi_0 = 2$, and $\xi_n = 1$ for $n \geq 1$.

3.7. Algorithm

1. Input (integer) n .

Input (double series) tools.

Input (array) $a_{old} = a_{00}$, (initial approximation, a_{00} with $n + 1$ dimension, are chosen so that the boundary conditions are satisfied).

2. $\hat{G} = (a_{old})$: $a_{new} = \hat{F}$ is a system of linear algebraic equations that has been solved and a_{new} is obtained.

Go to (2).

2.1 If $|a_{old} - a_{new}| < \text{tol}$ then $a_{new} = a$, break (the program is finished).

2.2 Else then $a_{old} \leftarrow a_{new}$.

3. Go to (2).

4. Main Results and Applications of Orthogonal Polynomials in Applied Science

In this section, will be applied the proposed methods to resolve some test problems of the non-homogeneous wave equation to get the approximate solution. We use two distinct error criteria to assess the accuracy of suggested methods, which were calculated by using Mathematica® 12. The norm L_{rms} is defined by:

$$L_{rms} = \left(\frac{\sum_{j=1}^N (u_{approx}(x_j, t_j) - u_{exact}(x_j, t_j))^2}{\sum_{j=1}^N (u_{exact}(x_j, t_j))^2} \right)^{1/2},$$

and the norm L_∞ which is defined by:

$$L_\infty = \max_{j=1,2,\dots,N} |u_{approx}(x_j, t_j) - u_{exact}(x_j, t_j)|.$$

where u_{approx} and u_{exact} represent the numerical and analytical solutions, respectively.

Example 1. Consider the following inhomogeneous wave problem:

$$\frac{\partial^2 u}{\partial t^2} = \frac{\partial^2 u}{\partial x^2} + 1, \quad t \geq 0, \quad 0 \leq x \leq 1. \quad (18)$$

with initial conditions:

$$u(x, 0) = x, \quad \frac{\partial u}{\partial t}(x, 0) = x, \quad (19)$$

and Dirichlet boundary conditions:

$$u(0, t) = \frac{t^2}{2}, \quad u(1, t) = 1 + t + \frac{t^2}{2}. \quad (20)$$

and the analytical solution is: $u(x, t) = x + xt + \frac{t^2}{2}$.

By applying the proposed method to resolve this problem with initial and boundary condition gives in **Equations 19**, and **20**, respectively. More specifically, by converting the unknown function $u(x, t)$ with partial derivatives in to linear equations system. Furthermore, assume that $n = 2$.

First: Applying the standard polynomials.

By inserting the **Equations 6**, and **7** into **Equations 18**, **19** and **20** the 2^{th} -order partial derivatives and conditions are converted into matrices, such that:

$$\begin{aligned} \psi(x).B.(A_S^2)^T.\phi(t) - \psi(x).A_S^2.B.\phi(t) &= 1, \\ [1 \ x \ x^2] \begin{bmatrix} b_{00} & b_{01} & b_{02} \\ b_{10} & b_{11} & b_{12} \\ b_{20} & b_{21} & b_{22} \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 2 & 0 & 0 \end{bmatrix} [1 \ t \ t^2]^T - \\ [1 \ x \ x^2] \begin{bmatrix} 0 & 0 & 2 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} b_{00} & b_{01} & b_{02} \\ b_{10} & b_{11} & b_{12} \\ b_{20} & b_{21} & b_{22} \end{bmatrix} [1 \ t \ t^2]^T &= 1, \\ 2b_{02} - 2b_{20} + 2b_{12}x - 2b_{21}t + 2b_{22}x^2 - 2b_{22}t^2 &= 1, \end{aligned} \quad (21)$$

For $n = 2$ the **Equation 6**, becomes as following:

$$\begin{aligned} u(x, t) &= [1 \ x \ x^2] \begin{bmatrix} b_{00} & b_{01} & b_{02} \\ b_{10} & b_{11} & b_{12} \\ b_{20} & b_{21} & b_{22} \end{bmatrix} [1 \ t \ t^2]^T, \\ u(x, t) &= b_{00} + b_{01}t + b_{02}t^2 + b_{10}x + b_{11}xt + b_{12}xt^2 + b_{20}x^2 + b_{21}x^2t + b_{22}x^2t^2, \end{aligned} \quad (22)$$

From **Equation 7**, the first derivatives of $u(x, t)$ with a respective t its:

$$\begin{aligned} \frac{\partial u(x, t)}{\partial t} &= \psi(x).B.(A_S)^T.\phi(t) = [1 \ x \ x^2] \begin{bmatrix} b_{00} & b_{01} & b_{02} \\ b_{10} & b_{11} & b_{12} \\ b_{20} & b_{21} & b_{22} \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 2 & 0 \end{bmatrix} [1 \ t \ t^2]^T, \\ &= b_{01} + 2b_{02}t + b_{11}x + 2b_{12}tx + b_{21}x^2 + 2b_{22}tx^2, \end{aligned} \quad (23)$$

By substituting the IC from **Equation 19**, when $t = 0$ in to **Equations 22**, and **23**, will be obtained:

$$u(x, 0) = \psi(x).A.\phi(0) = b_{00} + b_{10}x + b_{20}x^2 = x, \quad (24)$$

$$\frac{\partial u(x, 0)}{\partial t} = \psi(x).B.(A_S)^T.\phi(0) = b_{21}x^2 + b_{11}x + b_{01} = x, \quad (25)$$

Also, by inserting the BC from **Equation 20** when $x = 0, 1$ in to **Equation 22**, will be get:

$$u(0, t) = \psi(0).B.\phi(t) = b_{00} + b_{01}t + b_{02}t^2 = \frac{t^2}{2}, \quad (26)$$

$$u(1, t) = \psi(1).B.\phi(t),$$

$$u(1, t) = (b_{02} + b_{12} + b_{22})t^2 + (b_{01} + b_{11} + b_{21})t + (b_{00} + b_{10} + b_{20}) = 1 + t + \frac{t^2}{2}, \quad (27)$$

By resolve the linear algebraic system consisting of the **Equations 21**, **24**, **25**, **26**, and **27**, to find the values of b_{00} , b_{01} , b_{02} , b_{10} , b_{11} , b_{12} , b_{20} , b_{21} and b_{22} , using the Mathematica[®]12, we get:

$$B = \begin{bmatrix} b_{00} & b_{01} & b_{02} \\ b_{10} & b_{11} & b_{12} \\ b_{20} & b_{21} & b_{22} \end{bmatrix} = \begin{bmatrix} -1.1102 \times 10^{-15} & 0.2143 & 0.4643 \\ 1 & 0.3571 & 0.1071 \\ 3.3307 \times 10^{-16} & 0.4286 & -0.0714 \end{bmatrix}.$$

By replacing these values in **Equations 6** or **22**, therefore, the approximate solution using

standard polynomials (x^i) as follows:

$$u(x, t) \approx -1.1102 \times 10^{-15} + x + 3.3307 \times 10^{-16}x^2 \\ + t^2(0.4643 + 0.1071x - 0.0714x^2) + t(0.2143 + 0.3571x + 0.4286x^2).$$

Finally, by using the other proposed methods such as: Boubaker, Bernoulli, Chebyshev, Genocchi, and Legendre, the approximate solution for test example 1 as follows:

$$u(x, t) \approx x + t(-0.0455 + 1.13636x - 0.0909x^2) + t^2(0.5410 - 0.1410x + 0.1000x^2).$$

$$u(x, t) \approx x - 1.9660 \times 10^{-17}x^2 + t^2(0.5313 - 0.0938x + 0.0625x^2) \\ + t(0.0625 + 0.8125x + 0.125x^2).$$

$$u(x, t) \approx 1.7764 \times 10^{-15} + x - 1.1102 \times 10^{-15}x^2 \\ + t^2(0.6429 - 0.42857x + 0.2857x^2) \\ + t(0.2143 + 0.3571x + 0.4286x^2).$$

$$u(x, t) \approx -3.5527 \times 10^{-15} + x + 2.8866 \times 10^{-15}x^2 \\ + t(-0.7059 + 3.1176x - 1.4118x^2) \\ + t^2(2.3529 - 5.5588x + 3.7059x^2).$$

$$u(x, t) \approx 2.7756 \times 10^{-17} + x - 5.5511 \times 10^{-17}x^2 + t^2(0.5713 - 0.2139x + 0.1426x^2) \\ + t(0.2213 + 0.3361x + 0.4426x^2).$$

Table 1. The values of L_∞ and L_{rms} for example 1, when $n = 3, 4, \dots, 10$.

n	Norm	Standard Poly.	Boubaker Poly.	Bernoulli Poly.	Chebyshev Poly.	Genocchi Poly.	Legendre Poly.
3	L_∞	3.9389	6.3650	2.8073	4.1435	8.4519	2.8457
		$\times 10^{-4}$	$\times 10^{-4}$	$\times 10^{-4}$	$\times 10^{-3}$	$\times 10^{-3}$	$\times 10^{-3}$
	L_{rms}	1.3868	2.2434	9.8839	1.4605	3.1633	1.0040
		$\times 10^{-3}$	$\times 10^{-3}$	$\times 10^{-4}$	$\times 10^{-2}$	$\times 10^{-2}$	$\times 10^{-2}$
4	L_∞	2.8110	3.5392	5.2614	2.9529	1.2557	1.8394
		$\times 10^{-6}$	$\times 10^{-5}$	$\times 10^{-6}$	$\times 10^{-4}$	$\times 10^{-3}$	$\times 10^{-4}$
	L_{rms}	2.2211	2.8541	4.1481	2.3525	1.0075	1.4611
		$\times 10^{-5}$	$\times 10^{-4}$	$\times 10^{-5}$	$\times 10^{-3}$	$\times 10^{-2}$	$\times 10^{-3}$
5	L_∞	1.6801	2.0271	1.3806	3.1093	7.3779	3.9023
		$\times 10^{-8}$	$\times 10^{-6}$	$\times 10^{-7}$	$\times 10^{-5}$	$\times 10^{-5}$	$\times 10^{-6}$
	L_{rms}	1.9194	2.2173	1.5364	3.6629	8.9684	4.7726
		$\times 10^{-7}$	$\times 10^{-5}$	$\times 10^{-6}$	$\times 10^{-4}$	$\times 10^{-4}$	$\times 10^{-5}$
6	L_∞	7.0099	5.5441	3.4199	1.5229	5.0341	6.9510
		$\times 10^{-11}$	$\times 10^{-8}$	$\times 10^{-9}$	$\times 10^{-6}$	$\times 10^{-6}$	$\times 10^{-7}$
	L_{rms}	1.0776	8.5531	5.1635	2.3571	7.9422	1.0778
		$\times 10^{-9}$	$\times 10^{-7}$	$\times 10^{-8}$	$\times 10^{-5}$	$\times 10^{-5}$	$\times 10^{-5}$
7	L_∞	2.4247	1.4491	8.5369	1.2737	3.1418	2.7870
		$\times 10^{-13}$	$\times 10^{-9}$	$\times 10^{-11}$	$\times 10^{-9}$	$\times 10^{-7}$	$\times 10^{-8}$
	L_{rms}	4.5606	2.7474	1.5928	2.1049	6.1840	5.3380
		$\times 10^{-12}$	$\times 10^{-8}$	$\times 10^{-9}$	$\times 10^{-8}$	$\times 10^{-6}$	$\times 10^{-7}$
8	L_∞	8.8818	1.4573	1.0250	8.8493	9.4655	7.6410
		$\times 10^{-16}$	$\times 10^{-10}$	$\times 10^{-12}$	$\times 10^{-10}$	$\times 10^{-9}$	$\times 10^{-10}$
	L_{rms}	2.7022	3.3249	4.5441	4.0375	4.4433	3.4891
		$\times 10^{-14}$	$\times 10^{-9}$	$\times 10^{-11}$	$\times 10^{-8}$	$\times 10^{-7}$	$\times 10^{-8}$
9	L_∞	8.8818	2.0440	2.4425	1.1189	4.8860	5.7948
		$\times 10^{-16}$	$\times 10^{-11}$	$\times 10^{-14}$	$\times 10^{-10}$	$\times 10^{-9}$	$\times 10^{-11}$
	L_{rms}	3.2342	6.1451	6.4498	1.2092	2.6727	1.9853
		$\times 10^{-15}$	$\times 10^{-10}$	$\times 10^{-13}$	$\times 10^{-9}$	$\times 10^{-7}$	$\times 10^{-9}$
10	L_∞	4.4409	7.4571	3.5527	9.2631	1.3047	1.0119
		$\times 10^{-16}$	$\times 10^{-11}$	$\times 10^{-15}$	$\times 10^{-11}$	$\times 10^{-10}$	$\times 10^{-10}$
	L_{rms}	1.2699	4.9801	9.8615	1.0156	2.6093	1.2359
		$\times 10^{-14}$	$\times 10^{-9}$	$\times 10^{-13}$	$\times 10^{-8}$	$\times 10^{-7}$	$\times 10^{-8}$







 Norm $L_{\infty}, n = 3, 4, \dots, 10$.

 Norm $L_{rms}, n = 3, 4, \dots, 10$.

Figure 1. The plots logarithmic of L_{rms} and L_{∞} , for example 1, when $n = 3, 4, \dots, 10$.

In **Table 1** and **Figure 1**, it can be seen the values of the norm L_{rms} and L_{∞} . They depend on the value of n , which represents the degree of the polynomial. The higher the value of n gives the lower error. The best results were achieved using the standard polynomials when, $n = 10$, the values of L_{∞} and L_{rms} equals, 4.4409×10^{-16} , 1.2699×10^{-14} , respectively, followed by Bernoulli polynomials.


Figure 2. Plots the norm L_{∞} for example 1, when $n = 10$.


Exact solution



Approximate solution

Figure 3. Comparison between analytical and numerical solution for example 1 by the standard poly.

Example 2. Consider the inhomogeneous wave problem as follows:

$$\frac{\partial^2 u}{\partial t^2} = \frac{\partial^2 u}{\partial x^2} + (x^2 - 2) \cosh(t), \quad 0 \leq x \leq 1, t \geq 0, \quad (28)$$

with initial conditions:

$$u(x, 0) = x^2, \quad \frac{\partial u}{\partial t}(x, 0) = 0, \quad (29)$$

and Dirichlet boundary conditions:

$$u(0, t) = 0, \quad u(1, t) = \cosh(t). \quad (30)$$

and the analytical solution is: $u(x, t) = x^2 \cosh(t)$.

$$\psi(x).B.(A_n^{*2})^T.\phi(t) - \psi(x).A_n^{*2}.B.\phi(t) = (x^2 - 2) \cosh(t),$$

$$u(x, 0) = \psi(x).B.\phi(0) = x^2, \quad \frac{\partial u}{\partial t}(x, 0) = \psi(x).B.(A_n^*)^T.\phi(0) = 0,$$

$$u(0, t) = \psi(0).B.\phi(t) = 0, \quad u(1, t) = \psi(1).B.\phi(t) = \cosh(t),$$

such that A_n^* represented the operational matrices of all proposed methods.


Let us use the Mathematica[®]12 to solve the given test problem and determine the L_∞ and L_{rms} :

Table 2. The values of L_∞ and L_{rms} for example 2, when $n = 3, 4, \dots, 10$.


n	Norm	Standard Poly.	Boubaker Poly.	Bernoulli Poly.	Chebyshev Poly.	Genocchi Poly.	Legendre Poly.
3	L_∞	0.0188309	0.0188309	0.0436566	0.1571468	0.0430718	0.0689526
	L_{rms}	0.0704147	0.0704147	0.1338318	0.9958798	0.2414356	0.4054831
4	L_∞	1.3482 $\times 10^{-3}$	1.3482 $\times 10^{-3}$	4.0844 $\times 10^{-4}$	0.0355224	3.6705 $\times 10^{-3}$	0.0157769
	L_{rms}	0.0140385	0.0140385	4.5347 $\times 10^{-4}$	0.5368480	0.0560399	0.2322642
5	L_∞	9.3338 $\times 10^{-5}$	9.3338 $\times 10^{-5}$	4.4291 $\times 10^{-5}$	0.0168202	2.8721 $\times 10^{-4}$	5.5056 $\times 10^{-3}$
	L_{rms}	2.1050 $\times 10^{-3}$	2.1050 $\times 10^{-3}$	9.0735 $\times 10^{-5}$	0.3734015	0.0106238	0.1194339
6	L_∞	8.9516 $\times 10^{-6}$	8.9516 $\times 10^{-6}$	2.8350 $\times 10^{-6}$	4.7210 $\times 10^{-4}$	4.7884 $\times 10^{-5}$	1.9671 $\times 10^{-3}$
	L_{rms}	3.0845 $\times 10^{-4}$	3.0845 $\times 10^{-4}$	3.3746 $\times 10^{-6}$	0.0121814	1.7861 $\times 10^{-3}$	0.0570002
7	L_∞	4.3799 $\times 10^{-8}$	4.3799 $\times 10^{-8}$	1.0233 $\times 10^{-8}$	9.011 $\times 10^{-4}$	5.7300 $\times 10^{-6}$	7.0847 $\times 10^{-4}$
	L_{rms}	2.2095 $\times 10^{-6}$	2.2095 $\times 10^{-6}$	6.2162 $\times 10^{-7}$	0.1040413	2.6369 $\times 10^{-4}$	0.0253115
8	L_∞	5.3188 $\times 10^{-10}$	3.1343 $\times 10^{-10}$	1.8148 $\times 10^{-10}$	8.0279 $\times 10^{-10}$	9.0985 $\times 10^{-9}$	4.9988 $\times 10^{-10}$
	L_{rms}	1.0668 $\times 10^{-8}$	8.4181 $\times 10^{-9}$	1.1232 $\times 10^{-8}$	8.7810 $\times 10^{-8}$	8.3036 $\times 10^{-7}$	6.2008 $\times 10^{-8}$
9	L_∞	1.4315 $\times 10^{-11}$	1.5375 $\times 10^{-11}$	8.1493 $\times 10^{-12}$	4.0713 $\times 10^{-11}$	4.9242 $\times 10^{-9}$	3.1324 $\times 10^{-11}$
	L_{rms}	1.1569 $\times 10^{-9}$	1.3045 $\times 10^{-9}$	7.0331 $\times 10^{-10}$	8.4630 $\times 10^{-9}$	5.0793 $\times 10^{-7}$	8.4263 $\times 10^{-10}$
10	L_∞	1.2901 $\times 10^{-13}$	4.1190 $\times 10^{-11}$	1.7109 $\times 10^{-13}$	1.0473 $\times 10^{-10}$	1.6116 $\times 10^{-10}$	5.8388 $\times 10^{-11}$
	L_{rms}	1.0793 $\times 10^{-11}$	5.4818 $\times 10^{-9}$	3.0960 $\times 10^{-11}$	2.0187 $\times 10^{-8}$	4.9575 $\times 10^{-7}$	1.3722 $\times 10^{-8}$

The error norm and mean square error values are readily seen in **Table 2**, and **Figure 4**, which can be obtained by solving example 2 using Mathematica. Some methods gave close

results, but the Bernoulli polynomials gave better results when $n = 9$, such that L_∞, L_{rms} equals $8.1493 \times 10^{-12}, 7.0331 \times 10^{-10}$, respectively, but when $n = 10$ the Bernoulli and standard polynomials give the same results.




Norm $L_\infty, n = 3, 4, \dots, 10$.




Norm $L_{rms}, n = 3, 4, \dots, 10$.


Figure 4. The plots logarithmic of L_{rms} and L_∞ , for example 2, when $n = 3, 4, \dots, 10$.




Standard Polynomials




Boubaker Polynomials




Bernoulli Polynomials



Chebyshev Polynomials




Genocchi Polynomials




Legendre Polynomials

Figure 5. Plots the norm L_∞ for example 2, when $n = 10$.



Exact solution



Approximate solution

Figure 6. Comparison between analytical and numerical solution for example 2 by the Bernoulli poly.

Example 3. Consider the homogeneous wave problem as follows (7):

$$\frac{\partial^2 u}{\partial t^2} = \frac{\partial^2 u}{\partial x^2}, \quad 0 \leq x \leq 1, \quad t \geq 0. \quad (31)$$

with initial conditions:

$$u(x, 0) = x, \quad \frac{\partial u}{\partial t}(x, 0) = \cos(x), \quad (32)$$

and Neumann boundary conditions:

$$\frac{du}{dn}(0, t) = 1, \quad \frac{du}{dn}(1, t) = 1 - \sin(1) \sin(t). \quad (33)$$

and the analytical solution is: $u(x, t) = x + \cos(x) \sin(t)$.

By implementing the proposed methods will be get the following linear equations system:

$$\psi(x).B.(A_n^*)^T.\phi(t) - \psi(x).A_n^*.B.\phi(t) = 0,$$

$$u(x, 0) = \psi(x).A.\phi(0) = x, \quad \frac{\partial u}{\partial t}(x, 0) = \psi(x).B.(A_n^*)^T.\phi(0) = \cos(x),$$

$$u(0, t) = \psi(0).A_n^*.B.\phi(t) = 1, \quad \frac{du}{dn}(1, t) = \psi(1).A_n^*.B.\phi(t) = 1 - \sin(1) \sin(t).$$

Table 3. The values of L_∞ and L_{rms} for example 3, when $n = 3, 4, \dots, 10$.

n	Norm	Standard Poly.	Boubaker Poly.	Bernoulli Poly.	Chebyshev Poly.	Genocchi Poly.	Legendre Poly.
3	L_∞	0.026769	0.026769	8.1346×10^{-4}	0.422276	0.144075	0.165196
	L_{rms}	0.023984	0.023984	2.0740×10^{-3}	0.409013	0.204080	0.158717
4	L_∞	3.5980×10^{-3}	3.5980×10^{-3}	9.0742×10^{-4}	0.171641	0.025237	0.051985
	L_{rms}	3.6621×10^{-3}	3.6621×10^{-3}	3.6562×10^{-4}	0.231415	0.040730	0.069231
5	L_∞	2.4744×10^{-4}	2.4744×10^{-4}	1.9915×10^{-6}	1.0927×10^{-3}	5.3252×10^{-3}	0.015482
	L_{rms}	4.7965×10^{-4}	4.7965×10^{-4}	1.5268×10^{-5}	2.0580×10^{-3}	9.9614×10^{-3}	0.028927
6	L_∞	2.0079×10^{-5}	2.0079×10^{-5}	1.5870×10^{-6}	0.022490	3.7142×10^{-4}	4.5766×10^{-3}
	L_{rms}	5.5328×10^{-5}	5.5328×10^{-5}	8.5427×10^{-7}	0.056908	1.1897×10^{-3}	0.011584
7	L_∞	1.8479×10^{-6}	1.8268×10^{-6}	4.7259×10^{-9}	7.9157×10^{-7}	3.9991×10^{-5}	8.3372×10^{-6}
	L_{rms}	6.1196×10^{-6}	6.1167×10^{-6}	3.9008×10^{-8}	2.5910×10^{-6}	1.7296×10^{-4}	2.7798×10^{-5}
8	L_∞	1.5123×10^{-7}	2.8128×10^{-9}	2.1089×10^{-9}	5.7829×10^{-7}	9.2256×10^{-6}	2.5362×10^{-7}
	L_{rms}	6.3784×10^{-7}	3.7864×10^{-9}	8.9858×10^{-10}	2.4624×10^{-6}	5.1617×10^{-5}	1.0919×10^{-6}
9	L_∞	1.1721×10^{-8}	9.0889×10^{-9}	1.2325×10^{-11}	2.0718×10^{-8}	2.8711×10^{-7}	1.1747×10^{-10}
	L_{rms}	6.2317×10^{-8}	9.6519×10^{-9}	1.6392×10^{-11}	1.1274×10^{-7}	1.5240×10^{-6}	7.1236×10^{-10}
10	L_∞	8.9957×10^{-10}	7.1069×10^{-10}	2.6179×10^{-13}	2.3150×10^{-9}	1.0916×10^{-7}	1.4462×10^{-10}
	L_{rms}	5.8471×10^{-9}	5.8575×10^{-9}	8.0077×10^{-13}	1.4899×10^{-8}	7.0608×10^{-7}	9.8726×10^{-10}

From **Table 3.** and **Figure 7.**, the error norm and mean square error values are visible, as these values decrease as the value of n increases, which represents the degree of the polynomial. The best methods for approximation are the Bernoulli polynomials, such that when $n = 5$, the values of L_∞ and L_{rms} equals 8.1346×10^{-4} , 2.0740×10^{-3} ,

respectively, but when $n = 10$, it can be seen the values of L_∞ and L_{rms} gives 2.6179×10^{-13} , 8.0077×10^{-13} .




Figure 7. The plots logarithmic of L_{rms} and L_∞ , for example 3, when $n = 3, 4, \dots, 10$.




Figure 8. Plots the norm L_∞ for example 3, when $n = 10$.




Figure 9. Comparison between analytical and numerical solution for example 3 by the Bernoulli poly.

Example 4: Consider the following inhomogeneous wave problem (7):

$$\frac{\partial^2 u}{\partial t^2} = \frac{\partial^2 u}{\partial x^2} + 6t + 2x, \quad 0 \leq x \leq 1, t \geq 0, \quad (34)$$

with initial conditions:

$$u(x, 0) = 0, \quad \frac{\partial u}{\partial t}(x, 0) = \sin(x), \quad (35)$$

and Neumann boundary conditions:

$$\frac{du}{dn}(0, t) = t^2, \quad \frac{du}{dn}(1, t) = t^2 + \cos(1) \sin(t). \quad (36)$$

and the analytical solution is: $u(x, t) = t^3 + t^2x + \sin(x) \sin(t)$.

$$\psi(x).B.(A_n^{*2})^T.\phi(t) - \psi(x).A_n^{*2}.B.\phi(t) = 6t + 2x,$$




$$u(x, 0) = \psi(x).B.\phi(0) = 0, \quad \frac{\partial u}{\partial t}(x, 0) = \psi(x).B.(A_n^*)^T.\phi(0) = \sin(x),$$

$$\frac{du}{dn}(0, t) = \psi(0).A_n^*.B.\phi(t) = t^2, \quad \frac{du}{dn}(1, t) = \psi(1).A_n^*.B.\phi(t) = t^2 + \cos(1) \sin(t).$$


Table 4. The values error norm L_∞ and L_{rms} for example 4, when $n = 3, 4, \dots, 10$.

n	Norm	Standard Poly.	Boubaker Poly.	Bernoulli Poly.	Chebyshev Poly.	Genocchi Poly.	Legendre Poly.
3	L_∞	0.025288	0.025287	0.0010292	0.420795	0.420795	0.163715
	L_{rms}	0.021544	0.021544	0.0040845	0.409652	0.409652	0.157369
4	L_∞	3.6917 $\times 10^{-3}$	3.6917 $\times 10^{-3}$	1.0348 $\times 10^{-3}$	0.171735	6.7708 $\times 10^{-2}$	5.2079 $\times 10^{-2}$
	L_{rms}	3.7901 $\times 10^{-3}$	3.7901 $\times 10^{-3}$	2.7305 $\times 10^{-4}$	0.233360	9.1241 $\times 10^{-2}$	6.9889 $\times 10^{-2}$
5	L_∞	2.4244 $\times 10^{-4}$	2.4244 $\times 10^{-4}$	6.9924 $\times 10^{-6}$	8.9554 $\times 10^{-4}$	8.9811 $\times 10^{-3}$	1.5477 $\times 10^{-2}$
	L_{rms}	4.9382 $\times 10^{-4}$	4.9382 $\times 10^{-4}$	2.4738 $\times 10^{-5}$	1.7230 $\times 10^{-3}$	1.6941 $\times 10^{-2}$	2.9166 $\times 10^{-2}$
6	L_∞	1.9630 $\times 10^{-5}$	1.9630 $\times 10^{-5}$	2.3295 $\times 10^{-6}$	8.9054 $\times 10^{-3}$	1.0740 $\times 10^{-3}$	4.5761 $\times 10^{-3}$
	L_{rms}	5.5389 $\times 10^{-5}$	5.5389 $\times 10^{-5}$	7.7029 $\times 10^{-7}$	1.8313 $\times 10^{-2}$	2.7441 $\times 10^{-3}$	1.1675 $\times 10^{-2}$
7	L_∞	1.8584 $\times 10^{-6}$	1.8584 $\times 10^{-6}$	1.5232 $\times 10^{-8}$	3.7324 $\times 10^{-6}$	1.1798 $\times 10^{-4}$	8.5907 $\times 10^{-7}$
	L_{rms}	6.1401 $\times 10^{-6}$	6.1401 $\times 10^{-6}$	6.6218 $\times 10^{-8}$	1.2450 $\times 10^{-5}$	3.9714 $\times 10^{-4}$	2.7754 $\times 10^{-6}$
8	L_∞	1.5181 $\times 10^{-7}$	1.5182 $\times 10^{-7}$	2.6662 $\times 10^{-9}$	7.7151 $\times 10^{-7}$	4.1914 $\times 10^{-6}$	4.0559 $\times 10^{-7}$
	L_{rms}	6.4355 $\times 10^{-7}$	6.4355 $\times 10^{-7}$	1.3797 $\times 10^{-9}$	3.3122 $\times 10^{-6}$	1.8038 $\times 10^{-5}$	1.7364 $\times 10^{-6}$
9	L_∞	1.1711 $\times 10^{-8}$	1.2419 $\times 10^{-8}$	2.9055 $\times 10^{-11}$	3.4713 $\times 10^{-8}$	9.8285 $\times 10^{-7}$	1.3215 $\times 10^{-8}$
	L_{rms}	6.2850 $\times 10^{-8}$	5.9882 $\times 10^{-8}$	1.0129 $\times 10^{-11}$	1.8559 $\times 10^{-7}$	5.2592 $\times 10^{-6}$	7.0957 $\times 10^{-8}$
10	L_∞	8.9913 $\times 10^{-10}$	1.0768 $\times 10^{-9}$	2.9887 $\times 10^{-13}$	6.7981 $\times 10^{-10}$	8.1974 $\times 10^{-8}$	6.0933 $\times 10^{-10}$
	L_{rms}	5.8928 $\times 10^{-9}$	5.7708 $\times 10^{-9}$	8.2474 $\times 10^{-13}$	4.8477 $\times 10^{-9}$	7.1460 $\times 10^{-7}$	4.1418 $\times 10^{-9}$


From **Table 4**, and **Figure 10**, it can be seen the values of the L_{rms} and L_∞ . The best approximation method was the Bernoulli polynomial, where the value of the error norm was equal to 0.0010292 and the value of the mean square error was equal to 0.0040845 when $n = 3$, also when $n = 10$, the values of L_∞ , and L_{rms} are 2.9887×10^{-13} , 8.2474×10^{-13} , respectively.

Norm $L_\infty, n = 3, 4, \dots, 10$.Norm $L_{rms}, n = 3, 4, \dots, 10$.**Figure 10.** The plots logarithmic of L_{rms} and L_∞ , for example 4, when $n = 3, 4, \dots, 10$.


Standard Polynomials




Boubaker Polynomials




Bernoulli Polynomials




Chebyshev Polynomials




Genocchi Polynomials



Legendre Polynomials

Figure 11. Plots the norm L_∞ for example 4, when $n = 10$.

Exact solution



Approximate solution

Figure 12. Comparison between analytical and numerical solution for example 4 by the Bernoulli poly.**Example 5.** Consider the following homogeneous wave problem:

$$\frac{\partial^2 u}{\partial t^2} = \frac{\partial^2 u}{\partial x^2}, \quad t \geq 0, \quad 0 \leq x \leq 1, \quad (37)$$

with initial conditions:

$$u(x, 0) = x^2, \quad \frac{\partial u}{\partial t}(x, 0) = 0, \quad (38)$$

and mixed boundary conditions:

$$u(0, t) = t^2, \quad \frac{\partial u}{\partial n}(1, t) = 2. \quad (39)$$

and the analytical solution is: $u(x, t) = x^2 + t^2$.

$$\psi(x).B.(A_n^*)^T.\phi(t) - \psi(x).A_n^*.B.\phi(t) = 0,$$


$$u(x, 0) = \psi(x).B.\phi(0) = x^2, \quad \frac{\partial u}{\partial t}(x, 0) = \psi(x).B.(A_n^*)^T.\phi(0) = 0,$$

$$u(0, t) = \psi(0).B.\phi(t) = t^2, \quad \frac{\partial u}{\partial n}(1, t) = \psi(1).A_n^*.B.\phi(t) = 2.$$

Table 5. The values L_∞ and L_{rms} for example 5, when $n = 3, 4, \dots, 10$.

n	Norm	Standard Poly.	Boubaker Poly.	Bernoulli Poly.	Chebyshev Poly.	Genocchi Poly.	Legendre Poly.
3	L_∞	0.026367	0.02637	4.1199×10^{-4}	0.42187	0.421875	0.16479
	L_{rms}	0.028829	0.028829	4.5045×10^{-4}	0.46126	0.461264	0.18018
4	L_∞	2.6674×10^{-3}	2.6674×10^{-3}	1.0419×10^{-5}	0.17071	6.6684×10^{-2}	5.1055×10^{-2}
	L_{rms}	4.0600×10^{-3}	4.0600×10^{-3}	1.5859×10^{-5}	0.25984	0.10150	7.7711×10^{-2}
5	L_∞	2.4968×10^{-4}	2.4968×10^{-4}	2.4382×10^{-7}	6.3917×10^{-2}	8.9883×10^{-3}	1.5484×10^{-2}
	L_{rms}	5.2356×10^{-4}	5.2356×10^{-4}	5.1129×10^{-7}	0.13403	1.8848×10^{-2}	3.2469×10^{-2}
6	L_∞	2.1965×10^{-5}	2.1965×10^{-5}	5.3626×10^{-9}	5.9419×10^{-5}	1.0763×10^{-3}	1.7436×10^{-4}
	L_{rms}	6.2408×10^{-5}	6.2408×10^{-5}	1.5236×10^{-8}	1.6882×10^{-4}	3.0580×10^{-3}	4.9539×10^{-4}
7	L_∞	1.8433×10^{-6}	1.8433×10^{-6}	1.1251×10^{-10}	1.8693×10^{-6}	1.1797×10^{-4}	1.4866×10^{-5}
	L_{rms}	6.9147×10^{-6}	6.9147×10^{-6}	4.2211×10^{-10}	7.0124×10^{-6}	4.4254×10^{-4}	5.5767×10^{-5}
8	L_∞	1.4915×10^{-7}	4.0861×10^{-9}	2.2808×10^{-12}	4.3597×10^{-7}	1.2081×10^{-5}	6.3073×10^{-8}
	L_{rms}	7.1560×10^{-7}	1.7778×10^{-8}	1.0964×10^{-11}	2.0917×10^{-6}	5.7964×10^{-5}	3.0262×10^{-7}
9	L_∞	1.1727×10^{-8}	2.2072×10^{-9}	7.5362×10^{-13}	8.7881×10^{-9}	5.1274×10^{-7}	8.9391×10^{-8}
	L_{rms}	6.9919×10^{-8}	1.5722×10^{-9}	8.6578×10^{-12}	5.2527×10^{-8}	3.0578×10^{-6}	5.3297×10^{-7}
10	L_∞	9.0098×10^{-10}	4.2117×10^{-8}	1.7586×10^{-13}	7.9146×10^{-8}	9.7206×10^{-8}	1.0858×10^{-8}
	L_{rms}	6.5670×10^{-9}	4.0507×10^{-8}	2.6847×10^{-12}	4.4758×10^{-8}	7.1679×10^{-7}	2.9672×10^{-8}


The mean square error and the error norm values can be easily observed in **Table 5**, and **Figure 13**, which we obtained by solving example 5 using Mathematica. Some methods gave close results; except for the Bernoulli polynomial, which gave better results in general. The values of L_∞ , and L_{rms} when $n = 3$ can be observed to be equals to 4.1199×10^{-4} , 4.5045×10^{-4} and equals 1.7586×10^{-13} , 2.6847×10^{-12} , respectively, when $n = 10$.



Norm L_∞ , $n = 3, 4, \dots, 10$.

Norm L_{rms} , $n = 3, 4, \dots, 10$.

Figure 13. The plots logarithmic of L_{rms} and L_∞ , for example 5, when $n = 3, 4, \dots, 10$.



Standard Polynomials

Boubaker Polynomials


Bernoulli Polynomials

Chebyshev Polynomials

Genocchi Polynomials

Legendre Polynomials

Figure 14. Plots the norm L_∞ for example 5, when $n = 10$.



Exact solution

Approximate solution

Figure 15. Comparison between analytical and numerical solution for example 5 by the Bernoulli poly.

Example 6. Consider the following homogeneous wave problem:

$$\frac{\partial^2 u}{\partial t^2} = \frac{\partial^2 u}{\partial x^2}, \quad t \geq 0, \quad 0 \leq x \leq 1, \quad (40)$$

with initial conditions:

$$u(x, 0) = 0, \quad \frac{\partial u}{\partial t}(x, 0) = \cosh(x), \quad (41)$$

and mixed boundary conditions:

$$u(0, t) = \sinh(t), \quad \frac{\partial u}{\partial n}(1, t) = \sinh(1) \sinh(t), \quad (42)$$

and the analytical solution is: $u(x, t) = \cosh(x) \sinh(t)$.

$$\psi(x).B.(A_n^{*2})^T.\phi(t) - \psi(x).A_n^{*2}.B.\phi(t) = 0,$$

$$u(x, 0) = \psi(x).B.\phi(0) = 0, \quad \frac{\partial u}{\partial t}(x, 0) = \psi(x).B.(A_n^*)^T.\phi(0) = \cosh(x),$$




$$u(0, t) = \psi(0).B.\phi(t) = \sinh(t),$$

$$\frac{\partial u}{\partial n}(1, t) = \psi(1).A_n^*.B.\phi(t) = \sinh(1) \sinh(t).$$


Table 6. The values of L_∞ and L_{rms} for example 6, when $n = 3, 4, \dots, 10$.

n	Norm	Standard Poly.	Boubaker Poly.	Bernoulli Poly.	Chebyshev Poly.	Genocchi Poly.	Legendre Poly.
3	L_∞	0.02371	0.02371	0.00239	0.41921	0.41921	0.16213
	L_{rms}	2.3180 $\times 10^{-2}$	2.3180 $\times 10^{-2}$	8.0865 $\times 10^{-3}$	0.48249	0.48249	0.18379
4	L_∞	4.4663 $\times 10^{-3}$	4.4663 $\times 10^{-3}$	1.8093 $\times 10^{-3}$	0.17251	6.8483 $\times 10^{-2}$	5.2854 $\times 10^{-2}$
	L_{rms}	3.6227 $\times 10^{-3}$	3.6227 $\times 10^{-3}$	1.0568 $\times 10^{-3}$	0.27538	0.10710	8.1823 $\times 10^{-2}$
5	L_∞	2.5933 $\times 10^{-4}$	2.5933 $\times 10^{-4}$	9.8953 $\times 10^{-6}$	6.3926 $\times 10^{-2}$	8.9980 $\times 10^{-3}$	1.5493 $\times 10^{-2}$
	L_{rms}	5.0699 $\times 10^{-4}$	5.0698 $\times 10^{-4}$	5.1090 $\times 10^{-5}$	0.14239	1.9981 $\times 10^{-2}$	3.4457 $\times 10^{-2}$
6	L_∞	2.6267 $\times 10^{-5}$	2.6267 $\times 10^{-5}$	4.3072 $\times 10^{-6}$	1.5617 $\times 10^{-5}$	1.0806 $\times 10^{-3}$	1.0011 $\times 10^{-4}$
	L_{rms}	6.3034 $\times 10^{-5}$	6.3034 $\times 10^{-5}$	3.7037 $\times 10^{-6}$	3.0898 $\times 10^{-5}$	3.2466 $\times 10^{-3}$	3.1860 $\times 10^{-4}$
7	L_∞	1.8672 $\times 10^{-6}$	1.8672 $\times 10^{-6}$	2.4094 $\times 10^{-8}$	1.5815 $\times 10^{-6}$	1.1799 $\times 10^{-4}$	2.6682 $\times 10^{-5}$
	L_{rms}	7.2126 $\times 10^{-6}$	7.2126 $\times 10^{-6}$	1.3786 $\times 10^{-7}$	6.5372 $\times 10^{-6}$	4.7018 $\times 10^{-4}$	1.0614 $\times 10^{-4}$
8	L_∞	1.5414 $\times 10^{-7}$	1.5414 $\times 10^{-7}$	4.9978 $\times 10^{-9}$	1.0245 $\times 10^{-7}$	1.2086 $\times 10^{-5}$	3.2711 $\times 10^{-7}$
	L_{rms}	7.5332 $\times 10^{-7}$	7.5332 $\times 10^{-7}$	7.4583 $\times 10^{-9}$	5.5511 $\times 10^{-7}$	6.1594 $\times 10^{-5}$	1.7006 $\times 10^{-6}$
9	L_∞	1.1754 $\times 10^{-8}$	1.1754 $\times 10^{-8}$	2.6757 $\times 10^{-11}$	5.4474 $\times 10^{-8}$	6.7884 $\times 10^{-8}$	6.3814 $\times 10^{-8}$
	L_{rms}	7.4120 $\times 10^{-8}$	7.4123 $\times 10^{-8}$	1.8809 $\times 10^{-10}$	3.4480 $\times 10^{-7}$	4.2685 $\times 10^{-7}$	4.1004 $\times 10^{-7}$
10	L_∞	9.0452 $\times 10^{-10}$	9.0442 $\times 10^{-10}$	3.5627 $\times 10^{-12}$	2.8841 $\times 10^{-9}$	9.8860 $\times 10^{-8}$	1.5768 $\times 10^{-8}$
	L_{rms}	6.9709 $\times 10^{-9}$	6.9705 $\times 10^{-9}$	8.3315 $\times 10^{-12}$	9.7122 $\times 10^{-10}$	7.8001 $\times 10^{-7}$	1.2507 $\times 10^{-7}$


From **Table 6**, and **Figure 16**, the error norm and mean square error values are presented. The best approximation method was the Bernoulli polynomial, where the value of the error norm was equal to 3.5627×10^{-12} and the value of the mean square error was equal to 8.3315×10^{-12} when $n = 10$.

Norm $L_{\infty}, n = 3, 4, \dots, 10$.Norm $L_{rms}, n = 3, 4, \dots, 10$.**Figure 16.** The plots logarithmic of L_{rms} and L_{∞} , for example 6, when $n = 3, 4, \dots, 10$.


Standard Polynomials




Boubaker Polynomials




Bernoulli Polynomials




Chebyshev Polynomials




Genocchi Polynomials



Legendre Polynomials

Figure 17. Plots the norm L_{∞} for example 6, when $n = 10$.

Exact solution



Approximate solution

Figure 18. Comparison between analytical and numerical solution for example 6 by the Bernoulli poly.

5. Conclusion

In this work, the orthogonal polynomials and operational matrices based on standard polynomials were proposed and applied to resolve non-homogeneous wave problems. The approximate solutions were achieved and showed to be reliable and effective, even for low-order polynomials. Additionally, the mean square error L_{rms} and the norm L_{∞} were calculated to assess the accuracy, dependability, and validity of the approaches. The results

show that the recommended solutions have high accuracy and lower error rates. In all situations, the mean square error and error norm values drop as n grows. Furthermore, it is showed that the results of the L_∞ , and L_{rms} by the recommended methods Bernoulli and standard polynomials fell greatly in comparison to the other orthogonal polynomials. The standard polynomial was the best in the first example when $n = 5$ to 10, and the values of the L_∞ , and L_{rms} were equal to 4.4409×10^{-16} , 1.2699×10^{-14} , respectively, when $n = 10$, while the Bernoulli Polynomial was the best in the remaining examples as it gave the lowest error when $n = 3$ to 10.

Acknowledgment

The authors are very grateful to the referees for giving them insightful comments and ideas for enhancing the paper.

Conflict of Interest

The authors declare that they have no conflicts of interest.

Funding

There is no monetary support in preparing for the publication.

References

1. Wazwaz AM. Linear and nonlinear integral equations. Berlin: Springer; 2011.
2. He JH. Variational iteration method for delay differential equations. Commun Nonlinear Sci Numer Simul. 1997;2(4):235–236. [https://doi.org/10.1016/S1007-5704\(97\)90008-3](https://doi.org/10.1016/S1007-5704(97)90008-3)
3. He JH. Variational iteration method for autonomous ordinary differential systems. Appl Math Comput. 2000;114(2–3):115–123. [https://doi.org/10.1016/S0096-3003\(99\)00104-6](https://doi.org/10.1016/S0096-3003(99)00104-6)
4. He JH. Variational iteration method-Some recent results and new interpretations. J Comput Appl Math. 2007;207(1):3–17. <https://doi.org/10.1016/j.cam.2006.07.009>
5. Wazwaz AM. The variational iteration method: A powerful scheme for handling linear and nonlinear diffusion equations. Comput Math with Appl. 2007;54(7–8):933–939. <https://doi.org/10.1016/j.camwa.2006.12.038>
6. Wazwaz AM. The variational iteration method: A reliable analytic tool for solving linear and nonlinear wave equations. Comput Math with Appl. 2007;54(7–8):926–932. <https://doi.org/10.1016/j.camwa.2006.12.038>
7. Wazwaz AM. Partial differential equations and solitary waves theory. Springer Science & Business Media; 2010.
8. Mansour MB. Accurate computation of traveling wave solutions of some nonlinear diffusion equations. Wave Motion. 2007;44(3): 222-230. [doi:10.1016/j.wavemoti.2006.10.001](https://doi.org/10.1016/j.wavemoti.2006.10.001)
9. Biazar J, Ghazvini H. An analytical approximation to the solution of a wave equation by a variational iteration method. Appl Math Lett. 2008;21(8):780–5. <https://doi.org/10.1016/j.aml.2007.08.004>
10. Britt S, Tsynkov S, Turkel E. Numerical solution of the wave equation with variable wave speed on nonconforming domains by high-order difference potentials. J Comput Phys. 2018;354:26–42. <https://doi.org/10.1016/j.jcp.2017.10.049>
11. Cheniguel A. Numerical Method for Solving Wave Equation with Non Local Boundary Conditions. Proc Int MultiConference Eng Comput Sci. 2013;2:13–5.
12. Ullah, H., Islam, S., Dennis, L.C.C., Abdelhameed, T.N., Khan, I. and Fiza M. Approximate solution of two-dimensional nonlinear wave equation by optimal homotopy asymptotic method. Math Probl Eng. 2015;2015(1):380104. <https://doi.org/10.1155/2015/380104>
13. Abolarin OE. New Improved Variational Homotopy Perturbation Method for Bratu-Type Problems. Am J Comput Math. 2013;03(02):110–113. <https://doi.org/10.4236/ajcm.2013.32018>
14. Kumar S, Kumar A, Momani S, Aldhaifallah M, Nisar KS. Numerical solutions of nonlinear

- fractional model arising in the appearance of the stripe patterns in two-dimensional systems. *Adv Differ Equations*. 2019;2019(1):1-19. <https://doi.org/10.1186/s13662-019-2334-7>
- 15.15. Kumar S, Kumar A, Abbas S, Al Qurashi M, Baleanu D. A modified analytical approach with existence and uniqueness for fractional Cauchy reaction–diffusion equations. *Adv Differ Equations*. 2020;2020(1):28. <https://doi.org/10.1186/s13662-019-2488-3>
 - 16.16. Corrington MS. Solution of Differential and Integral Equations with Walsh Functions. *IEEE Trans Circuit Theory*. 1973;20(5):470–476.
 - 17.17. Sparis PD, Mouroutsos SG. The operational matrix of differentiation for orthogonal polynomial series. *Int J Control*. 1986;44(1):1–15. <https://doi.org/10.1080/00207178608933579>
 - 18.18. Kosunalp HY, G˘ulsu M. Operational Matrix by Hermite Polynomials for Solving Nonlinear Riccati Differential Equations. *Int J Math Comput Sci*. 2020;16(2):525–536.
 - 19.19. Bagherpoorfard M, Ghassabzade FA. Hermite Matrix Polynomial Collocation Method for Linear Complex Differential Equations and Some Comparisons. *J Appl Math Phys*. 2013;01(05):58–64. <http://dx.doi.org/10.4236/jamp.2013.15009>
 - 20.20. Isah A, Phang C. Operational matrix based on Genocchi polynomials for solution of delay differential equations. *Ain Shams Eng J*. 2018;9(4):2123–2128. <https://doi.org/10.1016/j.asej.2016.09.015>
 - 21.21. Abdelkawy MA, Taha TM. An operational matrix of fractional derivatives of laguerre polynomials. *Walailak J Sci Technol*. 2014;11(12):1041–1055. <http://dx.doi.org/10.14456/WJST.2014.59>
 - 22.22. Yousefi SA, Behroozifar M. Operational matrices of Bernstein polynomials and their applications. *Int J Syst Sci*. 2010;41(6):709–716. <https://doi.org/10.1080/00207720903154783>
 - 23.23. Yousefi SA, Behroozifar M, Dehghan M. The operational matrices of Bernstein polynomials for solving the parabolic equation subject to specification of the mass. *Journal of computational and applied mathematics*. 2011;235(17):5272-83. [doi:10.1016/j.cam.2011.05.038](https://doi.org/10.1016/j.cam.2011.05.038)
 - 24.24. Ibraheem GH, Al-Jawary MA. The operational matrix of Legendre polynomials for solving nonlinear thin film flow problems. *Alexandria Eng J*. 2020;59(5):4027–4033. <https://doi.org/10.1016/j.aej.2020.07.008>
 - 25.25. Sharma B, Kumar S, Paswan MK, and Mahato D. Chebyshev Operational Matrix Method for Lane-Emden Problem. *Nonlinear Eng*. 2019;8(1):1–9. <https://doi.org/10.1515/nleng-2017-0157>
 - 26.26. El-Gamel M E, Adel W, El-Azab MS. Collocation Method Based on Bernoulli Polynomial and Shifted Chebychev for Solving the Bratu Equation. *J Appl Comput Math*. 2018;07(03).
 - 27.27. Singh S, Patel VK, Singh VK, Tohidi E. Application of Bernoulli matrix method for solving two-dimensional hyperbolic telegraph equations with Dirichlet boundary conditions. *Comput Math with Appl*. 2018;75(7):2280–2294. <https://doi.org/10.1016/j.camwa.2017.12.003>
 - 28.28. Bencheikh A, Chiter L. Boubaker Operational Matrix Method for Fractional Emden-Fowler Problem. *J Sib Fed Univ - Math Phys*. 2023;16(3):357–369. <https://www.mathnet.ru/eng/jsfu1085>
 - 29.29. Turkyilmazoglu M. Effective computation of exact and analytic approximate solutions to singular nonlinear equations of Lane-Emden-Fowler type. *Appl Math Model*. 2013;37(14–15):7539–7548. <https://doi.org/10.1016/j.apm.2013.02.014>
 - 30.30. Turkyilmazoglu M. An effective approach for numerical solutions of high-order Fredholm integro-differential equations. *Appl Math Comput*. 2014;227:384–398. <https://doi.org/10.1016/j.amc.2013.10.079>
 - 31.31. Turkyilmazoglu M. Effective computation of solutions for nonlinear heat transfer problems in Fins. *J Heat Transfer*. 2014;136(9):1–7. <https://doi.org/10.1115/1.4027772>
 - 32.32. Turkyilmazoglu M. Solution of Initial and Boundary Value Problems by an Effective Accurate Method. *Int J Comput Methods*. 2017;14(6). <https://doi.org/10.1142/S0219876217500694>
 - 33.33. Turkyilmazoglu M. Parabolic Partial Differential Equations with Nonlocal Initial and Boundary Values. *Int J Comput Methods*. 2015;12(5):1–10. <https://doi.org/10.1142/S0219876215500243>
 - 34.34. Salih OM, AL-Jawary MA. Computational Methods for Solving Nonlinear Ordinary Differential Equations Arising in Engineering and Applied Sciences. *Iraqi J Sci*. 2023;64(8):4070–

4091. DOI: [10.24996/ijs.2023.64.8.30](https://doi.org/10.24996/ijs.2023.64.8.30)
- 35.35. Enadi MO, Al-Jawary MA. The operational matrices for Elliptic Partial Differential Equations with mixed boundary conditions. J Phys Conf Ser. 2024;2857(1):1–18. DOI [10.1088/1742-6596/2857/1/012027](https://doi.org/10.1088/1742-6596/2857/1/012027)
- 36.36. Enadi MO, AL-Jawary MA, Turkyilmazoglu M. The operational matrices methods for solving the Poisson equation with nonlocal boundary conditions. J Appl Math Comput. 2025;71(2):2385–2432. <https://doi.org/10.1007/s12190-024-02313-y>
- 37.37. Davaeifar S, Rashidinia J. Boubaker polynomials collocation approach for solving systems of nonlinear Volterra–Fredholm integral equations. J Taibah Univ Sci. 2017;11(6):1182–1199. <https://doi.org/10.1016/j.jtusci.2017.05.002>
- 38.38. Bhrawy AH, Tohidi E, Soleymani F. A new Bernoulli matrix method for solving high-order linear and nonlinear Fredholm integro-differential equations with piecewise intervals. Appl Math Comput. 2012;219(2):482–497. <https://doi.org/10.1016/j.amc.2012.06.020>
- 39.39. Toutounian F, Tohidi E. A new Bernoulli matrix method for solving second order linear partial differential equations with the convergence analysis. Appl Math Comput. 2013;223:298–310. <https://doi.org/10.1016/j.amc.2013.07.094>
- 40.40. Jung, C.Y., Liu, Z., Rafiq, A., Ali, F. and Kang SM. Solution of second order linear and nonlinear ordinary differential equations using Legendre operational matrix of differentiation. Int J Pure Appl Math. 2014;93(2):285–295. doi: <http://dx.doi.org/10.12732/ijpam.v93i2.12>
- 41.Doha EH, Bhrawy AH, Ezz-Eldien SS. A Chebyshev spectral method based on operational matrix for initial and boundary value problems of fractional order. Comput Math with Appl. 2011;62(5):2364–2373. <https://doi.org/10.1016/j.camwa.2011.07.024>.